



Implementing scalable face-to-face and digital interventions among forcibly displaced persons from Ukraine in Europe: protocol of The U-RISE Project

Karl J. Weinreich, Iryna Frankova, Natalie Maksymets, Corrado Barbui, Vitalii Klymchuk, Trudy Mooren, Emilia Olechno, Marianna Purgato, Marit Sijbrandij, Anke B. Witteveen & Els van der Ven

To cite this article: Karl J. Weinreich, Iryna Frankova, Natalie Maksymets, Corrado Barbui, Vitalii Klymchuk, Trudy Mooren, Emilia Olechno, Marianna Purgato, Marit Sijbrandij, Anke B. Witteveen & Els van der Ven (2025) Implementing scalable face-to-face and digital interventions among forcibly displaced persons from Ukraine in Europe: protocol of The U-RISE Project, *European Journal of Psychotraumatology*, 16:1, 2468117, DOI: [10.1080/20008066.2025.2468117](https://doi.org/10.1080/20008066.2025.2468117)

To link to this article: <https://doi.org/10.1080/20008066.2025.2468117>



© 2025 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 27 Feb 2025.



Submit your article to this journal [↗](#)



Article views: 1100



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)



STUDY PROTOCOL



Implementing scalable face-to-face and digital interventions among forcibly displaced persons from Ukraine in Europe: protocol of The U-RISE Project

Karl J. Weinreich ^a, Iryna Frankova ^{a,b}, Natalie Maksymets^a, Corrado Barbui ^d, Vitalii Klymchuk ^c, Trudy Mooren ^b, Emilia Olechno^a, Marianna Purgato ^d, Marit Sijbrandij ^a, Anke B. Witteveen ^a and Els van der Ven ^{a,b}

^aDepartment of Clinical, Neuro- and Developmental Psychology, World Health Organization Collaborating Center for Research and Dissemination of Psychological Interventions, Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands; ^bARQ National Psychotrauma Centre / Centrum 45, Oegstgeest, the Netherlands; ^cDepartment of Social Sciences, University of Luxembourg, Esch-sur-Alzette, Luxembourg; ^dWHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy

ABSTRACT

Background: The full-scale Russian invasion of Ukraine on 24 February 2022 has led to millions of forcibly displaced persons (FDPs) within Ukraine and other European countries. Due to war-related exposure and displacement adversities, this group is at significant risk of developing depression, anxiety, post-traumatic stress disorder, and other mental health problems. Systemic barriers, including insufficiently equipped mental health systems and language barriers, prevent FDPs from receiving adequate mental health and psychosocial support (MHPSS). Scalable interventions delivered in person by non-specialist helpers, or digitally, provide opportunities to scale up the MHPSS response.

Objectives: This paper aims to provide an overview of the 'Ukraine's displaced people in the EU: Reach out, Implement, Scale-up and Evaluate interventions promoting mental wellbeing' (U-RISE) project. U-RISE aims to improve the mental wellbeing of FDPs from Ukraine by establishing a network of Ukrainian mental health professionals, building sustainable capacity for provision and supporting implementation of scalable face-to-face and digital mental health interventions adapted to the specific needs of this population.

Method: We build capacity for and implement scalable face-to-face interventions, including Problem Management Plus, Self Help Plus, and Multi-family Approach, for FDPs from Ukraine in Poland, Slovakia and Romania. Digital interventions, including the Doing What Matters in Times of Stress digital guide and a Telegram-based chatbot 'Friend' using principles of Psychological First Aid, are being implemented in Europe and Ukraine. To monitor the population's mental wellbeing and impact of the interventions, qualitative needs assessments among mental health providers and FDPs, and quantitative assessments pre- and post-intervention are collected.

Conclusion: We provide a framework for the rapid implementation of face-to-face and digital interventions in countries that need to scale up their MHPSS in response to humanitarian or complex emergency crises.

Implementando intervenciones digitales y cara a cara escalables en personas forzosamente desplazadas de Ucrania en Europa: protocolo del Proyecto U-Rise

Antecedentes: La invasión rusa a gran escala en Ucrania el 24 de febrero de 2022 ha provocado que millones de personas se vean obligadas a desplazarse (FDPs en su sigla en inglés) dentro de Ucrania y hacia otros países europeos. Debido a la exposición relacionada con la guerra y las adversidades del desplazamiento, este grupo corre un riesgo significativo de desarrollar depresión, ansiedad, trastorno de estrés postraumático y otros problemas de salud mental. Las barreras sistémicas, incluidos los sistemas de salud mental insuficientemente equipados y las barreras lingüísticas, impiden que las FDP reciban un apoyo psicosocial y de salud mental adecuado (MHPSS en su sigla en inglés). Las intervenciones escalables realizadas en persona por interventores no especializados, o de forma digital, brindan oportunidades para ampliar la respuesta de MHPSS.

Objetivo: Este artículo tiene como objetivo proporcionar una visión general del proyecto 'Personas desplazadas de Ucrania en la UE: alcanzar, implementar, escalar y evaluar intervenciones que promueven el bienestar mental' (U-RISE en su sigla en inglés). U-RISE pretende mejorar el bienestar mental de las personas desplazadas de Ucrania mediante el

ARTICLE HISTORY

Received 31 October 2024
Revised 5 February 2025
Accepted 6 February 2025

KEYWORDS

Forcibly displaced persons; mental health and psychosocial support; scalable interventions; task-shifting; Ukraine

PALABRAS CLAVE

Intervenciones escalables; apoyo en salud mental y psicosocial; personas desplazadas forzosamente; cambio de tareas; Ucrania

KEY MESSAGES

- Scalable interventions delivered in person by non-specialists, or digitally, have the potential to scale up the mental health and psychosocial support in response to the mass displacement caused by the war in Ukraine.
- U-RISE aims to improve the mental wellbeing of forcibly displaced persons from Ukraine in Europe by building sustainable capacity and implementing multiple task-shifting programmes. These programmes include Self-Help Plus, Problem Management Plus and Multifamily Approach, as well as digital formats, including Doing What Matters in Times of Stress digital guide and a chatbot based on principles of Psychological First Aid.
- U-RISE provides a framework for the rapid implementation of scalable in-person and digital

establecimiento de una red de profesionales de la salud mental ucranianos, y crear una capacidad sostenible para la prestación y el apoyo a la implementación de intervenciones de salud mental escalables presenciales y digitales.

Método: Desarrollamos capacidad e implementamos intervenciones presenciales escalables, que incluyen Gestión de problemas plus, Autoayuda plus y Enfoque multifamiliar, para las FDPs de Ucrania en Polonia, Eslovaquia y Rumania. Se están implementando intervenciones digitales, que incluyen la guía digital Haciendo lo Que Importa en Tiempos de Estrés y un chatbot de Telegram 'Friend', que utiliza principios de Primeros auxilios psicológicos, en Europa y Ucrania. Para monitorear el bienestar mental de la población y el impacto de las intervenciones, se recopilan evaluaciones cualitativas de las necesidades entre los proveedores de salud mental y las FDP, y evaluaciones cuantitativas antes y después de la intervención.

Conclusión: Proporcionamos un marco de trabajo para la implementación rápida de intervenciones presenciales y digitales en países que necesitan ampliar su MHPSS en respuesta a crisis humanitarias o de emergencia complejas.

Abbreviations: DWM: Doing What Matters in Times of Stress; **ENACT:** Enhancing Assessment of Common Therapeutic Factors; **EQUIP:** Developed Ensuring Quality In Psychological Support Materials; **FDPs:** Forcibly Displaced Persons; **MFA:** Multi-Family Approach; **MHPSS:** Mental Health and Psychosocial Support; **NGO:** Non-Governmental Organisations; **PFA:** Psychological First Aid; **PM+:** Problem Management Plus; **PTSD:** Post-Traumatic Stress Disorder; **SH+:** Self-Help Plus; **ToH:** Training of Helpers; **ToT:** Training of Trainers; **UNHCR:** United Nations High Commissioner for Refugees; **U-RISE:** Ukraine's Displaced People in the EU: Reach Out, Implement, Scale-up and Evaluate Interventions Promoting Mental Wellbeing; **WHO:** World Health Organization

interventions underpinned by qualitative and quantitative research assessing the needs of various stakeholder groups and psychological outcomes pre- and post-intervention.

1. Introduction

On 24 February 2022, the Russian Federation invaded Ukraine, escalating the armed conflict that had been sweltering since Russia's annexation of Crimea eight years earlier, thereby initiating one of the largest humanitarian crises in Europe since World War II (Loft & Brien, 2023; United Nations High Commissioner for Refugees [UNHCR], 2024a). Since the invasion, more than three million individuals have been internally displaced, and almost 6 million individuals have been displaced into the European Union, the majority of whom are women and children (UN Women, 2023; UNHCR, 2024a). Besides stressors experienced during forced migration like violence, death, or other traumatic events, forcibly displaced persons (FDPs) from Ukraine face additional post-migration difficulties. These include loss of social status, cultural shock, language barriers, uncertainty related to temporary protection status, and reduced access to health care and psychosocial services (Frankova et al., 2024; UN Women, 2023).

These stressors occur in addition to other post-displacement stressors including the ongoing war, worries for the safety of loved ones back home due to growing documentation of war crimes, and fears concerning the potential use of nuclear weapons or damage to nuclear reactors (Independent International Commission of Inquiry on Ukraine, 2023; Morganstein et al., 2023; Porter & Haslam, 2005; UN Women, 2023). Furthermore, due to pan-Ukrainian mobilisation and conscription of the male population aged 25–60, men of fighting age are unable to leave Ukraine, resulting in women, children, and elderly, who are at higher risk of mental health problems

related to forced displacement, being overrepresented among FDPs from Ukraine (Jain et al., 2022; Li et al., 2016). Due to the measures taken to allow Ukrainians to move freely within the European Union, FDPs from Ukraine represent a highly mobile population that is hard to reach through conventional channels of mental health and psychosocial support (MHPSS).

Studies indicate that forced displacement has exacerbated pre-existing mental health problems among various groups of the Ukrainian population (Bai et al., 2022; Martsenkovskiy et al., 2022). In a cross-sectional survey among individuals who were internally displaced by the hostilities since 2014, prior to the current escalation, 22% reported symptoms of depression, 18% of anxiety, and 32% of post-traumatic stress disorder (PTSD; Roberts et al., 2017). Approximately 6 months after the Russian invasion, 25.9% of people still residing in Ukraine met self-reported criteria for PTSD (Karatzias et al., 2023). An online survey using quota sampling also reported high rates of psychological distress (52.7%), symptoms of anxiety (54.1%), and of depression (46.8%; Xu et al., 2023). These mental health problems are compounded by negative coping strategies (i.e. substance abuse) due to war exposure and are likely to increase in prevalence and severity as the war continues (Hyland et al., 2023; Karatzias et al., 2023).

2. Mental health services for FDPs from Ukraine in Europe

Currently, Slovakia hosts approximately 127,000, Romania 172,000, and Poland 980,000 FDPs from Ukraine, making their health services important first

points of contact for MHPSS (Eurostat, n.d.). Before the full-scale Russian invasion, mental health systems in these countries were already suffering from pre-existing challenges including high rates of unmet care needs, low availability of medical staff, underdiagnosis, stigma around mental health issues, and overreliance on inpatient services (Dlouhy, 2014; Winkler et al., 2017), which were only beginning to be addressed following the COVID-19 pandemic (Directorate-General for Health and Food Safety, 2023; Zadka & Olajossy, 2021). In most countries bordering Ukraine, such as Slovakia, Romania, and Poland, public mental health systems were overall lacking and local MHPSS services were consequently insufficiently equipped to meet the demand of individuals requiring timely mental health support (Organisation for Economic Co-operation and Development, 2023b, 2023c, 2023a; UNHCR, 2024b).

To reduce MHPSS service gaps, the World Health Organization (WHO) launched the Mental Health Gap Action Plan (WHO, 2008). Following the framework derived from this plan (WHO, 2016), the WHO developed several scalable MHPSS programmes that are applicable to broad population groups, transdiagnostic, and suitable for task-sharing. Task-sharing refers to the principle of delegating tasks originally performed by highly-trained specialists to less specialised workers, who receive additional training (van Ginneken et al., 2021). This allows for MHPSS interventions to be carried out by trained primary care workers, paraprofessionals or peer helpers under clinical supervision. Task-sharing may tackle issues like lack of qualified staff, waiting lists, and language barriers (Frankova et al., 2023). It is ideally implemented in stepped or collaborative care settings, where individuals with more severe mental health problems are referred to specialised services. In addition, guided and unguided digital psychological interventions have been developed to ensure delivery to a broad population through scalability, ease of access, and fast distribution (Cuijpers, 2023). Digital interventions can efficiently provide MHPSS to large groups of individuals and benefit from a lower barrier of entry for those who are hard to reach by face-to-face services because of stigma, residing in frontline zones, or high geographical mobility (Bear et al., 2024; Schueller et al., 2019).

3. The U-RISE Project

The present project protocol describes the ‘Ukraine’s displaced people in the EU: Reach out, Implement, Scale-up and Evaluate interventions promoting mental wellbeing’ (U-RISE) project, which was developed following a call by the European Union EU4Health Programme in May 2022 to contribute to establishing an adequate MHPSS response and addressing the

mental health needs of FDPs from Ukraine in bordering EU countries, specifically Poland, Romania and Slovakia (Figure 1). To this end, the U-RISE project has four main aims (Figure 2). First, U-RISE aims to mobilise and reinforce a European network of Ukrainian- and Russian-speaking mental health professionals. Difficulties in finding psychological support with sufficient cultural knowledge and language skills remain a challenge for displaced Ukrainians. Establishing such a network can help form a platform for sharing MHPSS resources and knowledge, connecting MHPSS providers with clients, and recruiting mental health professionals as trainers for scalable interventions. Second, U-RISE engages in capacity building by training mental health professionals to train peer helpers, thereby enabling non-governmental organisations (NGOs) to scale up MHPSS responses in the countries of implementation. In most Western European countries, MHPSS is provided by government-funded health systems. In countries bordering Ukraine, the MHPSS response is largely organised by humanitarian NGOs established through the interagency standing committee framework and coordinated via local technical working groups (Inter-Agency Standing Committee, 2007). Third, U-RISE aims to implement best practice face-to-face and digital interventions tailored to the needs of Ukrainian FDPs. Fourth, U-RISE aims to develop a roadmap of available specialised mental health services in Poland, Romania and Slovakia. This roadmap aims to inform MHPSS providers with an overview of available specialised support for Ukrainian FDPs, accelerate referral pathways, and enhance the efficiency of delivering care.

U-RISE is coordinated by Vrije Universiteit Amsterdam in collaboration with other academic institutions (University of Verona; University of Luxemburg), implementing NGOs (International Medical Corps Croatia (Poland); TENENET (Slovakia), Phoneo (Romania), ARQ National Psychotrauma Centre (Netherlands)), and guidance from WHO. A steering group composed of Ukrainian and European independent experts, and international partners such as the United Nations High Commissioner for Refugees (UNHCR), WHO Europe and WHO regional offices in Poland, Romania, and Slovakia, was established to provide external advice on the implementation, transferability, stakeholder needs and sustainability of U-RISE.

4. Ethical considerations

To safeguard ethical standards during implementation and evaluation, ethics approval for the study components was gathered from the Medical Ethics Review Committee of Amsterdam University Medical Centers (2023.0924), the Scientific and Ethical Review Board of



Figure 1. U-RISE implementation sites.

the Faculty of Behavioural and Movement Sciences of Vrije Universiteit Amsterdam (VCWE-2023-133) and the Institutional Review Board of the University of

Verona (12a/2023). To ensure the prioritisation of participant safety and well-being, referral pathways for participants in need of more intensive mental



Figure 2. U-RISE objectives overview.

health care were established in each implementation country prior to implementation. We used the Ensuring Quality In Psychological support materials (EQUIP; Marshall et al., 2010) to assess the skills of non-professional peer helpers and provided supervision to avoid beneficiaries would be exposed to unhelpful or potentially harmful behaviour. To ensure ethical treatment of participants, best practice principles were conveyed during trainings, and interventions were offered to all participants regardless of research participation.

5. Face to face interventions

The implementation of face-to-face interventions is carried out by the NGOs in Romania, Slovakia, and Poland. The U-RISE interventions include three face-to-face interventions for individuals (Problem Management Plus; PM+), groups (Self-Help Plus; SH+) and families (Multi-Family Approach; MFA).

5.1. Problem Management Plus

PM+ (WHO, 2020b) is a WHO-developed, scalable individual or group intervention (eight to 10 participants per group) aimed at reducing anxiety, depression, PTSD symptoms, and improving daily functioning for people affected by adversity. The intervention encompasses five weekly 90-minute sessions, delivered by peer-helpers under clinical supervision, in which participants focus on different strategies, including stress management, problem solving, behavioural activation, and social support strengthening skills (Dawson et al., 2015). Randomised controlled trials examining PM+ showed a reduction in symptoms of depression, anxiety, PTSD, and functional impairments in Pakistan (Rahman et al., 2016) and Kenya (Bryant et al., 2017). This evidence was extended to a trial with Syrian refugees in the Netherlands (de Graaff et al., 2023), even at six and 12-month follow-up (de Graaff et al., 2024; Schäfer et al., 2023). PM+ has been culturally adapted for Ukrainians and has since 2015 been implemented in non-occupied Donetsk and Luhansk regions for internally displaced persons and host communities (Goloktionova & Mukerjee, 2021). Ukrainian primary health professionals were trained as PM+ trainers and provided an eight-day training in the eastern regions of Ukraine to enable non-professional peer-helpers to deliver PM+ during the CHANGE project (Fuhr et al., 2021).

5.2. Self help plus

SH+ (WHO, 2021) is a WHO-developed, scalable, evidence-based group intervention (Augustinavicius

et al., 2023; Karyotaki et al., 2023; Tol et al., 2020). It aims to reduce distress and build resilience, and can be delivered to groups of up to 30 people. It is delivered by supervised, non-specialist facilitators who complete a short training and use pre-recorded audio, the illustrated guide *Doing What Matters in Times of Stress* (DWM), and scripted group exercises. Over five sessions lasting two hours each, participants learn strategies derived from Acceptance and Commitment Therapy, like grounding, unhooking, and accepting negative feelings. The intervention has been shown to reduce the risk of developing a mental disorder among refugees displaced to various European countries (Purgato et al., 2021) and Syrian refugees displaced to Turkey (Acarturk et al., 2022). In 2023, with support of the WHO Ukraine office, 90 trainers completed the Self-Help Plus stress management course. The trainers in Ukraine regularly scale the intervention by training facilitators to conduct group stress management sessions within state and civil society organisations in Ukraine (WHO, 2023).

5.3. Multi-Family Approach

MFA is a preventive intervention targeting multiple families sharing experiences of facing adversity (Mooren et al., 2020). It aims to improve parenting skills, attachment, inter- and intra-family support, and prevent generational consequences of stressors. MFA comprises of six sessions, lasting around 2.5 h each. Initial sessions focus on identifying stressors, and distinguishing between those that can and cannot be changed. The focus then shifts from the consequences of these stressors on individual physical and mental health, to interactions within the family. Lastly, the sessions focus on exploring and emphasising resources inside and outside the family. MFA is based on multi-family therapy (Asen & Scholz, 2010), which has been shown to improve family functioning (van Es et al., 2023). While the efficacy of MFA has not yet been empirically evaluated, it was found to be feasible and safe across other populations affected by conflict (Mooren et al., 2018). In 2023, over 40 professionals were trained in MFA.

6. Digital interventions

Digital interventions are being disseminated through partner websites in implementation countries and in Ukraine (e.g. by the Ukrainian mental health programme on the initiative of Olena Zelenska 'How are you?', or Ukrainian Ministry of Health), through leaflets, and word of mouth. Digital interventions are accessible worldwide. Meta-analyses suggest that guided and unguided digital MHPSS interventions are effective overall (Linardon et al., 2019), in low-

and middle-income countries (Fu et al., 2020), and as standalone interventions (Weisel et al., 2019).

6.1. Friend. First Aid Chatbot

The 'Friend' First Aid Chatbot is a digital preventive intervention based on principles of Psychological First Aid (PFA; WHO et al., 2011) that was adapted for Ukrainians dealing with stress due to ongoing hostilities (Frankova et al., 2022). The Chatbot was in development at Bogomolets National Medical University in Kyiv. It was launched in Ukrainian and English upon the full-scale Russian invasion of Ukraine in February 2022, being extended with a Russian language version in September 2022. The Chatbot is particularly suitable for hard-to-reach populations, as it can be accessed through the messenger service Telegram, used commonly in Ukraine. The Chatbot simulates a conversation with a specialist trained in PFA and provides recommendations and exercises for coping with acute stress (WHO et al., 2011). The Chatbot uses various scripts to adapt conversations based on participants' environments, distinguishing between life-threatening or safe environments (Lahutina et al., 2024). Although the Chatbot has not been empirically evaluated in a randomised controlled trial, its content is based on PFA, which offers practical and emotional support to reduce stress and promote coping in the immediate aftermath of a crisis or disaster and has been found to decrease acute distress and PTSD symptoms (Figuroa et al., 2022; WHO et al., 2011) and improved overall functioning following acute traumatic events (McCart et al., 2020). Feasibility study results have additionally shown a medium to large effect of the Chatbot on stress levels (Lahutina et al., 2024).

6.2. Doing what matters in times of stress

The DWM digital smartphone intervention is a self-help stress management guide for coping with adversity, adapted from WHO's illustrated self-help guide from the SH+ programme (Mediavilla et al., 2023; WHO, 2020a). DWM is freely available in Ukrainian and Russian via a website (dwmatters.eu). The intervention consists of five modules covering mindfulness-based strategies for managing stress: grounding, unhooking, acting on one's values, being kind, and making room. DWM uses pre-recorded audio exercises to support identifying facilitators and barriers, as well as stress triggers for participants. Guided DWM has been found to be effective at reducing symptoms of depression and anxiety among health workers affected by distress (Mediavilla et al., 2023) and refugees and migrants (Purgato et al. in press; Riello et al., 2021). Further, DWM has shown

promising results as a standalone app (Fu et al., 2020; Linardon et al., 2019; Weisel et al., 2019).

7. Intervention stratification and selection

U-RISE interventions are provided in a stratified manner (Figure 3). After contacting an NGO, FDPs receive a link to the Chatbot and are assigned to a peer helper. FDPs complete screenings of around 20 min with their peer helper, during which the risk of self-harm, other harm, suicide, and signs of psychosis are assessed. Should any of these risks be present, the individual is immediately referred to specialised mental health services. Subsequently, peer helpers decide which interventions are optimal, depending on the preference and perceived benefit of an individual or group intervention, and whether the individual experiences challenges in parenting. During this process, peer helpers can discuss individual cases in supervision sessions, and refer individuals to specialised care.

8. Quality assessment

To ensure the quality of trainings and care provision, the competency of peer helpers is assessed using the WHO-developed EQUIP materials (Marshall et al., 2010). EQUIP consists of freely available competency assessment tools and e-learning courses to support training institutions and NGOs, to train and supervise their workforce to deliver effective psychological support in humanitarian and development settings. EQUIP encompasses the Enhancing Assessment of Common Therapeutic Factors (ENACT) competency evaluation tool. ENACT assesses the mental health and psychosocial support skills for non-specialist and specialist providers across cultures, contexts and intervention types (Kohrt et al., 2020).

9. Capacity building

Building capacity for face-to-face interventions enables local NGOs in Romania, Slovakia, and Poland to scale-up their MHPSS and train hundreds of professionals and peer-helpers. Although a high turnover of staff and volunteers is expected, there are signals that a substantial proportion of Ukrainian refugees intend to settle in the EU (van Tubergen et al., 2024). Integrating this group into the MHPSS workforce is critical to address the unmet need for mental health support among FDPs from Ukraine.

For PM+ and SH+, a training pyramid consisting of trainings of trainers (ToTs) and subsequent trainings of peer-helpers (ToH), provides a structured framework for training a large number of trainers and peer-helpers and maximises the capacity for providing mental health support. Capacity building for PM+ and SH+ focuses on two sites in Poland, two




Figure 3. Intake and intervention trajectory of individuals.

in Romania and two in Slovakia. ToTs are provided to trainers, predominantly Ukrainian mental health professionals, employed or recruited by the NGOs. Trainers will train peer-helpers in ToH to deliver the interventions to FDPs seeking mental health support under supervision of a trainer.

In each country (Poland, Romania, and Slovakia) one ToT for PM + and SH + is organised, during which 15 trainers are trained. Afterwards, the aim is that each trainer trains five to seven peer-helpers, resulting in an estimated 100 peer helpers per country. Trainers provide 15–30 h of supervision to peer helpers. Aiming for five individual PM + interventions per helper, an estimated 400 FDPs per country and 1250 FDPs overall

receive an intervention. MFA is trained in two locations, Poland and Slovakia, followed by monthly online supervision sessions, during which facilitators are encouraged to set-up and offer family group sessions, and can discuss ongoing proceedings. For the digital interventions (Chatbot and DWM), no helpers need to be trained.

The U-RISE project aims to support the continuous scaling up of MHPSS. Monthly webinars for MHPSS providers are organised, alongside information dissemination through social media and traditional media. These webinars aim to connect Ukrainian mental health practitioners with international experts and facilitate knowledge and skill transfer in psychotrauma management amidst the challenges of the ongoing

conflict. These sessions offer 75 min of interactive communication, engaging experts from high-, middle-, and low-income countries. The online meetings are open for live participants and are available as recordings in English and Ukrainian. Further, training materials and guidance regarding implementation are offered to NGOs to disseminate evidence on the effectiveness of low-intensity interventions for FDPs from Ukraine, as well as guidance regarding implementation.

10. Data collection and evaluation

Qualitative data is collected through semi-structured key informant interviews with Ukrainian MHPSS providers, Ukrainian FDPs, and Ukrainian FDPs who received an intervention. Interviews last between 30 and 60 min each and rely on open-ended questions aimed at establishing the perceived needs of FDPs from Ukraine regarding the future scalability of MHPSS, factors currently impacting their mental health, and feedback for future adaptations. Interviews are recorded and transcribed through videoconferencing or audio recording software. Key themes and topics are identified using hybrid deductive and inductive analysis using ATLAS.ti and NVivo.

Quantitative data on mental health outcomes is collected at pre- and post-intervention among FDPs receiving PM+, SH+, Chatbot, or DWM. Mental health outcomes include depressive symptoms, anxiety symptoms, and health disability, assessed using the Patient Health Questionnaire (PHQ-9; Kroenke et al., 2001), Generalized Anxiety Disorder Scale (GAD-7; Spitzer et al., 2006), PTSD checklist (PCL-5; Blevins et al., 2015), and WHO Disability Assessment Schedule (WHODAS 2.0; WHO, 2010) respectively. Parental warmth and responsiveness are assessed using 14 items from the Parenting Scale (El-Khani et al., 2021). As there is not yet evidence on scalable MHPSS in Ukrainians residing in Europe, we aim to document the overall levels of mental distress in FDPs from Ukraine and any post-intervention changes.

11. Discussion

Europe is currently experiencing its most significant war since World War II, which has forcibly displaced more than 6 million persons from Ukraine into Europe. FDPs from Ukraine are at elevated risk for developing mental health complaints such as depression, anxiety, and PTSD due to exposure to war-related adversities. Many countries receiving high volumes of FDPs have insufficient capacity to adequately support their MHPSS needs, creating an urgent mental health care gap. The present paper describes the U-RISE project protocol, which aims to build a strong network of Ukrainian mental health

professionals, to contribute to capacity building for a scalable mental health response and to implement face-to-face and digital interventions. U-RISE promotes the use and dissemination of scalable interventions and addresses the MHPSS service gap in a real-world setting in a practical and time-efficient manner, thereby providing a framework for the quick implementation of face-to-face and digital interventions across multiple countries. The project evaluates the implementation of interventions in three bordering countries and assesses the needs of various stakeholder groups as well as mental health outcomes of FDPs receiving the interventions. With its complementary approach on capacity building, implementation and research components, including the structured evaluation of the implementation process, U-RISE can serve as an example for future projects focused on scaling up MHPSS in response to humanitarian crises.

Acknowledgements

We are very grateful to all participants in the U-RISE project, including master trainers, trainers, peer helpers and beneficiaries. Consortium members who have been key to the execution of the project, not listed as co-authors, include Gabriela Lupea, Federica Patania, Kateryna Khyzhniak, Emrah Kucukozkan, Anna Zubachova, Katarina Čavojská, Elena Kopcová, Martin, Mnahoncak, Zhannat Mukhamdiyeva, Olena Koval, Monica Bartucz, Nosheen Kashif and Michelle Engels.










Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by the European Union EU4H programme [grant number 101101495]. The opinions expressed in this paper are those of the authors and do not necessarily represent the decisions, policies, or views of the organisations they serve or the funder. The funders are not liable for any use that may be made of the information contained therein. Data summarised in the panel resulted from studies that had approval from the relevant ethics boards.

ORCID

Karl J. Weinreich  <https://orcid.org/0009-0005-7115-3631>
 Iryna Frankova  <https://orcid.org/0000-0001-9371-6849>
 Corrado Barbui  <https://orcid.org/0000-0003-1073-9282>
 Vitalii Klymchuk  <https://orcid.org/0000-0001-7898-5530>
 Trudy Mooren  <https://orcid.org/0000-0003-0819-8640>
 Marianna Purgato  <https://orcid.org/0000-0002-3783-8195>
 Marit Sijbrandij  <https://orcid.org/0000-0001-5430-9810>
 Anke B. Witteveen  <https://orcid.org/0000-0002-9636-7522>
 Els van der Ven  <http://orcid.org/0000-0001-5103-7186>

References

- Acarturk, C., Uygun, E., Ilkkursun, Z., Carswell, K., Tedeschi, F., Batu, M., Eskici, S., Kurt, G., Anttila, M., Au, T., Baumgartner, J., Churchill, R., Cuijpers, P., Becker, T., Koesters, M., Lantta, T., Nosè, M., Ostuzzi, G., Popa, M., ... Barbui, C. (2022). Effectiveness of a WHO self-help psychological intervention for preventing mental disorders among Syrian refugees in Turkey: A randomized controlled trial. *World Psychiatry*, 21(1), 88–95. <https://doi.org/10.1002/wps.20939>
- Asen, E., & Scholz, M. (2010). *Multi-family therapy: Concepts and techniques*. Routledge. <https://www.routledge.com/Multi-Family-Therapy-Concepts-and-Techniques/Asen-Scholz/p/book/9780415557818>.
- Augustinavicius, J., Purgato, M., Tedeschi, F., Musci, R., Leku, M. R., Carswell, K., Lakin, D., van Ommeren, M., Cuijpers, P., Sijbrandij, M., Karyotaki, E., Tol, W. A., & Barbui, C. (2023). Prevention and promotion effects of Self Help Plus: Secondary analysis of cluster randomised controlled trial data among South Sudanese refugee women in Uganda. *BMJ Open*, 13(9), e048043. <https://doi.org/10.1136/BMJOPEN-2020-048043>
- Bai, W., Cai, H., Sha, S., Ng, C. H., Javed, A., Mari, J., Su, Z., Shinfuku, N., & Xiang, Y. T. (2022). A joint international collaboration to address the inevitable mental health crisis in Ukraine. *Nature Medicine*, 28(6), 1103–1104. <https://doi.org/10.1038/s41591-022-01828-w>
- Bear, H. A., Ayala Nunes, L., Ramos, G., Manchanda, T., Fernandes, B., Chabursky, S., Walper, S., Watkins, E., & Fazel, M. (2024). The acceptability, engagement, and feasibility of mental health apps for marginalized and underserved young people: Systematic review and qualitative study. *Journal of Medical Internet Research*, 26, e48964. <https://doi.org/10.2196/48964>
- Blevins, C. A., Weathers, F. W., Davis, M. T., Witte, T. K., & Domino, J. L. (2015). The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and initial psychometric evaluation. *Journal of Traumatic Stress*, 28(6), 489–498. <https://doi.org/10.1002/jts.22059>
- Bryant, R. A., Schafer, A., Dawson, K. S., Anjuri, D., Mulili, C., Ndogoni, L., Koyiet, P., Sijbrandij, M., Ulate, J., Shehadeh, M. H., Hadzi-Pavlovic, D., & van Ommeren, M. (2017). Effectiveness of a brief behavioural intervention on psychological distress among women with a history of gender-based violence in urban Kenya: A randomised clinical trial. *PLOS Medicine*, 1–20. <https://doi.org/10.1371/journal.pmed.1002371>
- Cuijpers, P. (2023). The Dodo Bird and the need for scalable interventions in global mental health—a commentary on the 25th anniversary of Wampold et al. (1997). *Psychotherapy Research*, 33(4), 524–526. <https://doi.org/10.1080/10503307.2022.2138792>
- Dawson, K. S., Bryant, R. A., Harper, M., Kuowei Tay, A., Rahman, A., Schafer, A., & van Ommeren, M. (2015). Problem Management Plus (PM+): A WHO transdiagnostic psychological intervention for common mental health problems. *World Psychiatry*, 14(3), 354–357. <https://doi.org/10.1002/WPS.20255>
- de Graaff, A. M., Cuijpers, P., Elsayy, M., Hunaidy, S., Kieft, B., Gorgis, N., Twisk, J. W. R., Zakarian, Y., Bouman, T. K., Lommen, M. J. J., Acarturk, C., Bryant, R., McDaid, D., Morina, N., Park, A.-L., Ventevogel, P., & Sijbrandij, M. (2024). The effectiveness of Problem Management Plus at 1-year follow-up for Syrian refugees in a high-income setting. *Epidemiology and Psychiatric Sciences*, 33, e50. <https://doi.org/10.1017/S2045796024000519>
- de Graaff, A. M., Cuijpers, P., Twisk, J. W. R., Kieft, B., Hunaidy, S., Elsayy, M., Gorgis, N., Bouman, T. K., Lommen, M. J. J., Acarturk, C., Bryant, R., Burchert, S., Dawson, K. S., Fuhr, D. C., Hansen, P., Jordans, M., Knaevelsrud, C., McDaid, D., Morina, N., ... Sijbrandij, M. (2023). Peer-provided psychological intervention for Syrian refugees: Results of a randomised controlled trial on the effectiveness of Problem Management Plus. *BMJ Mental Health*, 26(1), 1–10. <https://doi.org/10.1136/bmjment-2022-300637>
- Directorate-General for Health and Food Safety. (2023). *Communication from the commission to the European Parliament, the council, the European Economic and Social Committee and the committee of the regions*. European Commission. https://health.ec.europa.eu/publications/comprehensive-approach-mental-health_en
- Dlouhy, M. (2014). Mental health policy in Eastern Europe: A comparative analysis of seven mental health systems. *BMC Health Services Research*, 14(1), 1–8. <https://doi.org/10.1186/1472-6963-14-42>
- El-Khani, A., Cartwright, K., Maalouf, W., Haar, K., Zehra, N., Çokamay-Yilmaz, G., & Calam, R. (2021). Enhancing teaching recovery techniques (Trt) with parenting skills: Rct of trt + parenting with trauma-affected Syrian refugees in Lebanon utilising remote training with implications for insecure contexts and covid-19. *International Journal of Environmental Research and Public Health*, 18(16), 1–20. <https://doi.org/10.3390/ijerph18168652>
- Eurostat. (n.d.). *Beneficiaries of temporary protection at the end of the month by citizenship, age and sex - monthly data*. Retrieved March 9, 2024, from https://ec.europa.eu/eurostat/databrowser/view/migr_asytpsm/default/table?lang=en.
- Figueroa, R. A., Cortés, P. F., Marín, H., Vergés, A., Gillibrand, R., & Repetto, P. (2022). The ABCDE psychological first aid intervention decreases early PTSD symptoms but does not prevent it: Results of a randomized-controlled trial. *European Journal of Psychotraumatology*, 13(1), 1–14. <https://doi.org/10.1080/20008198.2022.2031829>
- Frankova, I., Bahmad, M. L., Goloktionova, G., Suvalo, O., Khyzhniak, K., & Power, T. (2024). *Mental health and psychosocial support in Ukraine: Coping, help-seeking and health systems strengthening in times of war*. <https://arq.org/en/publications/mhps-ukraine>.
- Frankova, I., Klymchuck, V., Nickerson, A., Sijbrandij, M., & Amstadter, A. B. (2023). A summary of the 38th Annual International Society for traumatic stress studies presidential panel: How the traumatic stress community can assist individuals affected by the war in Ukraine. *Journal of Traumatic Stress*, 36(4), 682–690. <https://doi.org/10.1002/JTS.22945>
- Frankova, I., Vermetten, E., Shalev, A. Y., Sijbrandij, M., Holmes, E. A., Ursano, R., Schmidt, U., & Zohar, J. (2022). Digital psychological first aid for Ukraine. *The Lancet Psychiatry*, 9(7), e33. [https://doi.org/10.1016/S2215-0366\(22\)00147-X](https://doi.org/10.1016/S2215-0366(22)00147-X)
- Fu, Z., Burger, H., Arjadi, R., & Bockting, C. L. H. (2020). Effectiveness of digital psychological interventions for mental health problems in low-income and middle-income countries: A systematic review and meta-analysis. *The Lancet Psychiatry*, 7(10), 851–864. [https://doi.org/10.1016/S2215-0366\(20\)30256-X](https://doi.org/10.1016/S2215-0366(20)30256-X)

- Fuhr, D., Bogdanov, S., Tol, W., Nadkarni, A., & Roberts, B. (2021). Problem Management Plus and alcohol (PM + A): A new intervention to address alcohol misuse and psychological distress among conflict-affected populations. *Intervention, 19*(1), 141–143. https://doi.org/10.4103/INTV.INTV_27_20
- Goloktionova, A., & Mukerjee, M. (2021). Bringing Problem Management Plus to Ukraine: Reflections on the past and ways forward. *Intervention, 19*(1), 131–135. https://doi.org/10.4103/INTV.INTV_42_20
- Hyland, P., Vallières, F., Shevlin, M., Karatzias, T., Ben-Ezra, M., McElroy, E., Vang, M. L., Lorberg, B., & Martsenkovskiy, D. (2023). Psychological consequences of war in Ukraine: Assessing changes in mental health among Ukrainian parents. *Psychological Medicine, 53*(15), 7466–7468. <https://doi.org/10.1017/S0033291723000818>
- Independent International Commission of Inquiry on Ukraine. (2023). *Report of the independent international commission of inquiry on Ukraine*. UN Human Rights Council. <https://www.ohchr.org/en/hr-bodies/hrc/iicuhr-ukraine/index>.
- Inter-Agency Standing Committee. (2007). *IASC guidelines on mental health and psychosocial support in emergency settings*. <https://interagencystandingcommittee.org/iasc-task-force-mental-health-and-psychosocial-support-emergency-settings/iasc-guidelines-mental-health-and-psychosocial-support-emergency-settings-2007>.
- Jain, N., Prasad, S., Czárth, Z. C., Chodnekar, S. Y., Mohan, S., Savchenko, E., Panag, D. S., Tanasov, A., Betka, M. M., Platos, E., Świątek, D., Krygowska, A. M., Rozani, S., Srivastava, M., Evangelou, K., Gristina, K. L., Bordeniuc, A., Akbari, A. R., Jain, S., ... Reinis, A. (2022). War psychiatry: Identifying and managing the neuropsychiatric consequences of armed conflicts. *Journal of Primary Care & Community Health, 13*, 1–11. <https://doi.org/10.1177/21501319221106625>
- Karatzias, T., Shevlin, M., Ben-Ezra, M., McElroy, E., Redican, E., Vang, M. L., Cloitre, M., Ho, G. W. K., Lorberg, B., Martsenkovskiy, D., & Hyland, P. (2023). War exposure, posttraumatic stress disorder, and complex posttraumatic stress disorder among parents living in Ukraine during the Russian war. *Acta Psychiatrica Scandinavica, 147*(3), 276–285. <https://doi.org/10.1111/acps.13529>
- Karyotaki, E., Sijbrandij, M., Purgato, M., Acarturk, C., Lakin, D., Bailey, D., Peckham, E., Uygun, E., Tedeschi, F., Wancata, J., Augustinavicius, J., Carswell, K., Välimäki, M., van Ommeren, M., Koesters, M., Popa, M., Leku, M. R., Anttila, M., Churchill, R., ... Barbui, C. (2023). Self-Help Plus for refugees and asylum seekers: An individual participant data meta-analysis. *BMJ Mental Health, 26*(1), 1–8. <https://doi.org/10.1136/bmjment-2023-300672>
- Kohrt, B. A., Schafer, A., Willhoite, A., van't Hof, E., Pedersen, G. A., Watts, S., Ottman, K., Carswell, K., & van Ommeren, M. (2020). Ensuring Quality in Psychological Support (WHO EQUIP): Developing a competent global workforce. *World Psychiatry, 19*(1), 115. <https://doi.org/10.1002/WPS.20704>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine, 16*(9), 606–613. <https://doi.org/10.1046/J.1525-1497.2001.016009606.X>
- Lahutina, S., Frankova, I., Gruen, M., Vermetten, E., Zohar, J., Spitschan, M., & Bajbouj, M. (2024). A digital self-help tool to promote mental well-being for Ukrainians affected by war - assessing predictors of stress. *Neuroscience Applied, 3*, 104089. <https://doi.org/10.1016/J.NSA.2024.104089>
- Li, S. S. Y., Liddell, B. J., & Nickerson, A. (2016). The relationship between post-migration stress and psychological disorders in refugees and asylum seekers. *Current Psychiatry Reports, 18*(9), 1–9. <https://doi.org/10.1007/S11920-016-0723-0>
- Linardon, J., Cuijpers, P., Carlbring, P., Messer, M., & Fuller-Tyszkiewicz, M. (2019). The efficacy of app-supported smartphone interventions for mental health problems: A meta-analysis of randomized controlled trials. *World Psychiatry, 18*(3), 325–336. <https://doi.org/10.1002/wps.20673>
- Loft, P., & Brien, P. (2023). *Ukraine: UK aid and humanitarian situation 2022-23*. House of Commons Library. <https://commonslibrary.parliament.uk/research-briefings/cbp-9467/>.
- Marshall, J. C., Smart, J., & Horton, R. M. (2010). The design and validation of equip: An instrument to assess inquiry-based instruction. *International Journal of Science and Mathematics Education, 8*(2), 299–321. <https://doi.org/10.1007/S10763-009-9174-Y>
- Martsenkovskiy, D., Martsenkovsky, I., Martsenkovska, I., & Lorberg, B. (2022). The Ukrainian paediatric mental health system: Challenges and opportunities from the Russo-Ukrainian war. *The Lancet Psychiatry, 9*(7), 533–535. [https://doi.org/10.1016/S2215-0366\(22\)00148-1](https://doi.org/10.1016/S2215-0366(22)00148-1)
- McCart, M. R., Chapman, J. E., Zajac, K., & Rheingold, A. A. (2020). Community-based randomized controlled trial of psychological first aid with crime victims. *Journal of Consulting and Clinical Psychology, 88*(8), 681–695. <https://doi.org/10.1037/ccp0000588>
- Mediavilla, R., Felez-Nobrega, M., McGreevy, K. R., Monistrol-Mula, A., Bravo-Ortiz, M. F., Bayón, C., Giné-Vázquez, I., Villaescusa, R., Muñoz-Sanjose, A., Aguilar-Ortiz, S., Figueiredo, N., Nicaise, P., Park, A. la, Petri-Romão, P., Purgato, M., Witteveen, A. B., Underhill, J., Barbui, C., Bryant, R., ... Ayuso-Mateos, J. L. (2023). Effectiveness of a mental health stepped-care programme for healthcare workers with psychological distress in crisis settings: a multicentre randomised controlled trial. *BMJ Mental Health, 26*(1), 1–8. <https://doi.org/10.1136/BMJMENT-2023-300697>
- Mooren, T., Bala, J., & Rabaia, Y. (2020). A family-centered approach to working with refugee children and adolescents. *Child, Adolescent and Family Refugee Mental Health: A Global Perspective, 243–255*. https://doi.org/10.1007/978-3-030-45278-0_15
- Mooren, T., Rabaia, Y., Mitwalli, S., Reiffers, R., Koenen, R., & De Man, M. (2018). The multi-family approach to facilitate a family support network for Palestinian parents of children with a disability: A descriptive study. *The Lancet, 391*, S49. [https://doi.org/10.1016/S0140-6736\(18\)30415-X](https://doi.org/10.1016/S0140-6736(18)30415-X)
- Morganstein, J. C., Bromet, E. J., & Shigemura, J. (2023). The neuropsychiatric aftermath of exposure to weapons of mass destruction: Applying historical lessons to protect health during the war in Ukraine. *Psychological Medicine, 53*(11), 5353–5355. <https://doi.org/10.1017/S0033291722002872>
- Organisation for Economic Co-operation and Development. (2023a). *Poland: Country Health Profile 2023*. https://www.oecd.org/en/publications/poland-country-health-profile-2023_f597c810-en.html.
- Organisation for Economic Co-operation and Development. (2023b). *Romania: Country Health Profile*

2023. https://www.oecd.org/en/publications/romania-country-health-profile-2023_f478769b-en.html.
- Organisation for Economic Co-operation and Development. (2023c). *Slovakia: Country Health Profile 2023*. https://www.oecd.org/en/publications/slovakia-country-health-profile-2023_98f25a9b-en.html.
- Porter, M., & Haslam, N. (2005). Pre-displacement and post-displacement factors associated with mental health of refugees and internally displaced persons A meta-analysis. *JAMA*, 294(5), 602–612. <https://doi.org/10.1001/jama.294.5.602>
- Purgato, M., Carswell, K., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Bajbouj, M., Baumgartner, J., Biondi, M., Churchill, R., Cuijpers, P., Koesters, M., Gastaldon, C., Ilkkursun, Z., Lantta, T., Nosè, M., Ostuzzi, G., Papola, D., Popa, M., ... Barbui, C. (2021). Effectiveness of self-help plus in preventing mental disorders in refugees and asylum seekers in Western Europe: A multinational randomized controlled trial. *Psychotherapy and Psychosomatics*, 90(6), 403–414. <https://doi.org/10.1159/000517504>
- Rahman, A., Hamdani, S. U., Awan, N. R., Bryant, R. A., Dawson, K. S., Khan, M. F., Azeemi, M. M. U. H., Akhtar, P., Nazir, H., Chiumento, A., Sijbrandij, M., Wang, D., Farooq, S., & Van Ommeren, M. (2016). Effect of a multicomponent behavioral intervention in adults impaired by psychological distress in a conflict-affected area of Pakistan: A randomized clinical trial. *JAMA*, 316(24), 2609–2617. <https://doi.org/10.1001/jama.2016.17165>
- Riello, M., Purgato, M., Bove, C., Tedeschi, F., MacTaggart, D., Barbui, C., & Rusconi, E. (2021). Effectiveness of self-help plus (SH+) in reducing anxiety and post-traumatic symptomatology among care home workers during the COVID-19 pandemic: A randomized controlled trial. *Royal Society Open Science*, 8(11), 1–23. <https://doi.org/10.1098/RSOS.210219>
- Roberts, B., Makhshvili, N., & Javakhishvili, J. (2017). *Hidden burdens of conflict issues of mental health and access to services among internally displaced persons in Ukraine*. International Alert. <https://www.international-alert.org/publications/hidden-burdens-of-conflict/>.
- Schäfer, S. K., Thomas, L. M., Lindner, S., & Lieb, K. (2023). World Health Organization's low-intensity psychosocial interventions: A systematic review and meta-analysis of the effects of Problem Management Plus and step-by-step. *World Psychiatry*, 22(3), 449–462. <https://doi.org/10.1002/WPS.21129>
- Schueller, S. M., Hunter, J. F., Figueroa, C., & Aguilera, A. (2019). Use of digital mental health for marginalized and underserved populations. *Current Treatment Options in Psychiatry*, 6(3), 243–255. <https://doi.org/10.1007/s40501-019-00181-z>
- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097. <https://doi.org/10.1001/ARCHINTE.166.10.1092>
- Tol, W. A., Leku, M. R., Lakin, D. P., Carswell, K., Augustinavicius, J., Adaku, A., Au, T. M., Brown, F. L., Bryant, R. A., Garcia-Moreno, C., Musci, R. J., Ventevogel, P., White, R. G., & van Ommeren, M. (2020). Guided self-help to reduce psychological distress in South Sudanese female refugees in Uganda: A cluster randomised trial. *The Lancet Global Health*, 8(2), e254–e263. [https://doi.org/10.1016/S2214-109X\(19\)30504-2](https://doi.org/10.1016/S2214-109X(19)30504-2)
- UN Women. (2023). *Gender brief: Livelihoods and access to work of refugees from Ukraine in neighboring host countries*. https://eca.unwomen.org/sites/default/files/2023-09/livelihoods_brief_en.pdf.
- United Nations High Commissioner for Refugees. (2024a). *Ukraine Situation Flash Update #74*. <https://data.unhcr.org/en/documents/details/112054>.
- United Nations High Commissioner for Refugees. (2024b). *Summary of the humanitarian needs and response plan and the regional refugee response plan*. <https://reliefweb.int/report/ukraine/ukraine-summary-humanitarian-needs-and-response-plan-and-regional-refugee-response-plan-january-2024>.
- van Es, C. M., El Khoury, B., van Dis, E. A. M., te Brake, H., van Ee, E., Boelen, P. A., & Mooren, T. (2023). The effect of multiple family therapy on mental health problems and family functioning: A systematic review and meta-analysis. *Family Process*, 62(2), 499–514. <https://doi.org/10.1111/famp.12876>
- van Ginneken, N., Chin, W. Y., Lim, Y. C., Ussif, A., Singh, R., Shahmalak, U., Purgato, M., Rojas-García, A., Uphoff, E., McMullen, S., Foss, H. S., Thapa Pachya, A., Rashidian, L., Borghesani, A., Henschke, N., Chong, L. Y., & Lewin, S. (2021). Primary-level worker interventions for the care of people living with mental disorders and distress in low- and middle-income countries. *Cochrane Database of Systematic Reviews*, 2021(8), 1–642. <https://doi.org/10.1002/14651858.CD009149.pub3>
- van Tubergen, F., Wachter, G. G., Kosyakova, Y., & Kogan, I. (2024). Return intentions among Ukrainian refugees in Europe: A cross-national study. *International Migration*, 62(5), 181–198. <https://doi.org/10.1111/IMIG.13291>
- Weisel, K. K., Fuhrmann, L. M., Berking, M., Baumeister, H., Cuijpers, P., & Ebert, D. D. (2019). Standalone smartphone apps for mental health—a systematic review and meta-analysis. *Npj Digital Medicine*, 2(1), 1–10. <https://doi.org/10.1038/s41746-019-0188-8>
- Winkler, P., Krupchanka, D., Roberts, T., Kondratova, L., Machů, V., Höschl, C., Sartorius, N., van Voren, R., Aizberg, O., Bitter, I., Cerga-Pashoja, A., Deljkovic, A., Fanaj, N., Germanavicius, A., Hinkov, H., Hovsepian, A., Ismayilov, F. N., Ivezic, S. S., Jarema, M., ... Thornicroft, G. (2017). A blind spot on the global mental health map: A scoping review of 25 years' development of mental health care for people with severe mental illnesses in central and Eastern Europe. *The Lancet Psychiatry*, 4(8), 634–642. [https://doi.org/10.1016/S2215-0366\(17\)30135-9](https://doi.org/10.1016/S2215-0366(17)30135-9)
- World Health Organization. (2008). *mhGAP: Mental health gap action programme: Scaling up care for mental, neurological and substance use disorders*. <https://www.who.int/teams/mental-health-and-substance-use/treatment-care/mental-health-gap-action-programme>.
- World Health Organization. (2010). *Measuring health and disability manual for WHO Disability Assessment Schedule WHODAS 2.0* (T. B. Üstün, N. Kostanjek, S. Chatterij, & J. Rehm, Eds.). WHO Press. [https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-\(whodas-2.0\)](https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-(whodas-2.0)).
- World Health Organization. (2016). *mhGAP intervention guide for mental, neurological and substance use disorders in non-specialized health settings: Vol. Version 2.0*. <https://www.who.int/publications/i/item/9789241549790>.
- World Health Organization. (2020a). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications-detail-redirect/9789240003927>.
- World Health Organization. (2020b). *Group Problem Management Plus (Group PM+): Group psychological*

- help for adults impaired by distress in communities exposed to adversity (generic field-trial version 1.0).* <https://www.who.int/publications/i/item/9789240008106>.
- World Health Organization. (2021). *Self Help Plus (SH+): A group-based stress management course for adults. Generic field-trial version 1.0.* <https://www.who.int/publications/i/item/9789240035119>.
- World Health Organization. (2023). *WHO response to the Ukraine crisis: March 2023 Bulletin.* WHO European Region. <https://www.who.int/europe/publications/i/item/WHO-EURO-2023-6172-45937-69323>.
- World Health Organization, War Trauma Foundation, & World Vision International. (2011). *Psychological first aid: Guide for field workers.* World Health Organization. <https://www.who.int/publications/i/item/9789241548205>.
- Xu, W., Pavlova, I., Chen, X., Petrytsa, P., Graf-Vlachy, L., & Zhang, S. X. (2023). Mental health symptoms and coping strategies among Ukrainians during the Russia-Ukraine war in March 2022. *International Journal of Social Psychiatry*, 69(4), 957–966. <https://doi.org/10.1177/00207640221143919>
- Zadka, Ł, & Olajosy, M. (2021). COVID-19 Pandemic as the beginning of a golden era for telepsychiatry in Poland's healthcare system. *Frontiers in Psychiatry*, 11, 1–4. <https://doi.org/10.3389/fpsyt.2020.555559>