




BMJ Open Access and use of general and mental health services before and during the COVID-19 pandemic: a systematic review and meta-analysis

Camilla Sculco ¹, Beatrice Bano,^{1,2} Eleonora Prina ², Federico Tedeschi,^{2,3} Monica Bianca Bartucz,² Corrado Barbui,^{2,3} Marianna Purgato ^{2,3}, Emiliano Albanese¹

To cite: Sculco C, Bano B, Prina E, *et al.* Access and use of general and mental health services before and during the COVID-19 pandemic: a systematic review and meta-analysis. *BMJ Open* 2025;**15**:e091342. doi:10.1136/bmjopen-2024-091342

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-091342>).

MP and EA contributed equally.

MP and EA are joint senior authors.

Received 18 July 2024

Accepted 10 February 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

¹Institute of Public Health, Università della Svizzera italiana, Lugano, Ticino, Switzerland

²Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy

³Cochrane Global Mental Health, University of Verona, Verona, Italy

Correspondence to

Camilla Sculco;
camilla.sculco@usi.ch

ABSTRACT

Objectives To quantify access to health services during the COVID-19 pandemic and measure the change in use between the prepandemic and the pandemic periods in a population with assessment of psychological distress or diagnosis of mental disorders.

Data sources We developed and piloted a search syntax and adapted it to enter the following databases from 1 January 2020 to 31 March 2023: PubMed/MEDLINE, PsycINFO, Web of Science, Epistemonikos and the WHO International Clinical Trials Registry Platform. We reran the searches from the end of the original search to 3 December 2024.

Design We systematically screened titles, abstracts and full texts of retrieved records.

Eligibility criteria We included observational studies on any populations and regions, covering health services such as doctor visits, hospital admissions, diagnostic examinations, pharmaceutical therapies and mental health (MH) services. Only studies using validated scales to assess psychological distress or mental disorders as defined in the Diagnostic and Statistical Manual of Mental Disorders were included.

Data extraction and synthesis We extracted data using a purposefully designed form and evaluated the studies' quality with the Newcastle-Ottawa Scale. We measured the incidence rate (IR) of access to health services and the IR ratio (IRR) between the prepandemic and the pandemic periods. We calculated contacts days and catchment areas in the different periods. We used the random effects DerSimonian-Laird inverse-variance model and calculated heterogeneity with statistics I^2 and τ^2 . We computed pooled IR and pooled IRR and tested the hypothesis of no variation (IRR=1).

Results We retrieved 10 014 records and examined the full text of 580 articles. We included 136 primary studies of which 44 were meta-analysed. The IR of access to services during the pandemic was 2.59 contact months per 10 000 inhabitants (IR=2.592; 95% CI: 1.301 to 5.164). We observed a reduction of 28.5% in the use of services with negligible differences by age group and type of services (IRR=0.715; 95% CI: 0.651 to 0.785). We observed significant differences in effect sizes across studies ($\tau^2=5.44$; $p<0.001$ and $\tau^2=0.090$; $p<0.001$).

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Our review provides novel evidence on the association between mental health (MH) and the use of health services, focusing on populations with assessment of psychological distress or diagnosis of mental disorders. It includes studies from high-income and low- and middle-income countries.
- ⇒ The study's internal validity is strengthened by advanced statistical analyses, standardised procedures and rigorous quality assessments, ensuring reliability and minimising bias.
- ⇒ The included studies were conducted during different pandemic waves and across various countries, adding complexity to the consistency of outcomes.
- ⇒ The search was limited to five databases, language restrictions were applied; it was completed in March 2023, indicating the need for an update by the end of 2025.

Conclusion By considering MH, our study provides consolidated evidence and quantifies the reduction in the use of health services during the COVID-19 pandemic.
PROSPERO registration number CRD42023403778.

INTRODUCTION

The use of health services depends on the synergy between service characteristics such as availability, accessibility and acceptability and the health-seeking behaviours of the population.¹ The COVID-19 pandemic provides a unique opportunity to quantify variations in the use of health services driven by the interplay of an unprecedented reorganisation of services^{2–5} and a concomitant increase in mental health (MH) needs within the population. Globally, the reallocation of healthcare resources towards COVID-19 treatment coupled with the need to prevent in-hospital viral transmission significantly reduced the capacity for both elective and emergency procedures.⁶ Numerous studies



have documented substantial declines in emergency department visits,⁷ hospital admissions for acute medical conditions, including cardiovascular and coronary heart diseases,^{8 9} and presentations for MH conditions,^{10–15} along with reduced outpatient care contacts for physical health conditions.^{16–19} Besides, the pandemic has exerted a considerable and complex impact on the population's MH.^{2 20–27} Studies conducted since the outbreak reported rising rates of Common Mental Disorders (CMDs) such as depression, anxiety and stress within the general population. These increases have been attributed to several factors, including uncertainty about the pandemic's progression, the risk of infection, quarantine directives, social isolation, inactivity and income loss.^{28 29} However, the direction of the association between MH and the use of health services during the pandemic remains uncertain, as high levels of psychological distress within the population may be linked to both positive and negative attitudes towards seeking care.³⁰ In this context, a systematic synthesis of the evidence on the impact of the pandemic on the use of health services while also accounting for MH is lacking. The evidence is dispersed across a vast array of publications, and available studies have heterogeneity in outcome measures, settings and methodological approaches.²¹ Our goal was to consolidate evidence on the impact of the COVID-19 pandemic on the use of health services, providing a sound overview of available studies. Moreover, we aimed to fill the gap concerning the unclear association between MH and the use of health services by contemplating two approaches: focusing on a population with assessment of psychological distress or diagnosis of mental disorders and differentiating the use of general health services (ie, non-MH services) and MH services. The aims of this systematic review were (O1) to quantify the use of both general health and MH services during the COVID-19 pandemic and (O2) to assess the change in use of both general health and MH services between the prepandemic and the pandemic periods. We hypothesised that the pandemic has had an impact on the use of health services in several ways. On one hand, the adverse effects of COVID-19 on MH might have caused greater demand for care provision³¹; on the other hand, fear of infection, lack of outpatient care and non-availability of treatment in hospital settings might have prevented patients from seeking and using care.³² Understanding how the population used services is crucial to prioritise short- and long-term healthcare strategies and mitigating adverse physical and MH consequences of public health (PH) emergencies.³³

METHODS

The protocol for this review was registered in the International Prospective Register of Systematic Reviews, registration number CRD42023403778. We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for preparing the manuscript.

Data source

We developed and piloted a search syntax (online supplemental material) and adapted it to enter the following databases from 1 January 2020 to 31 March 2023: PubMed/MEDLINE, PsycINFO, Web of Science, Epistemonikos and the WHO International Clinical Trials Registry Platform. We hand-searched the reference lists of potentially relevant studies and considered additional studies suggested by the authors. We reran the searches with the exact syntax across the databases with specifications of dates between the end of the original search and 3 December 2024.

Study selection

We used the Population, Exposure, Outcome and Study Design framework to define the study selection. Inclusion criteria were: (a) observational studies; (b) study populations of any age, gender and subpopulation groups, such as minorities, patients with pre-existing psychological distress or mental disorders defined in the Diagnostic and Statistical Manual of Mental Disorders, 5th edition, Text Revision³⁴ and other pre-existing clinical conditions; (c) setting: studies carried out in any geographic region, as classified by the World Bank's 2022–2023 country income levels ranging from low- and middle-income to high-income³⁵; (d) type of health service: doctor visits, hospital admissions, diagnostic examinations and screenings, pharmaceutical therapies (intakes) and MH services (both inpatient and outpatient visits carried out by MH professionals such as psychiatrists, psychologists and psychotherapists); (e) studies that applied validated scales, which either directly assessed psychological distress or diagnosed mental disorders. In studies reporting assessment of psychological distress or diagnosis of mental disorders, we examined the use of both general health and MH services. In studies assessing psychological distress or diagnosis of mental disorders, we exclusively considered the use of MH services and (f) outcome measures reported by caregivers or other healthcare professionals instead of patients. Exclusion criteria were: (a) studies conducted in selected professional groups in which the MH burden of the pandemic was work-related; (b) studies reporting non-quantitative data, or information on referrals but not actual admissions, or prescriptions without information on actual intakes; (c) studies reporting generic questions about patients' status (eg, How do you feel today?) for the assessment of psychological distress; (d) studies retracted from publication and (e) studies published in languages other than English, Italian, French. After retrieving the records by bibliographic database search, we used the reference management databases EndNote³⁶ and Rayyan³⁷ to manage the screening, store and organise the references and remove duplicates. Three independent reviewers (CS, BB and EP) screened titles, abstracts and then the full texts of retrieved records. Disagreements were resolved by consensus with senior authors (MP and EA). The per cent agreement among reviewers prior to full-text screening was 82.15%.

Data extraction and synthesis

CS, BB and EP independently extracted data on primary studies' characteristics and results using a purposefully designed extraction form implemented in Microsoft Excel. We contacted the authors to supply data in predefined, analysis-ready formats when the data were incomplete or unclear. We performed periodic checks to ensure the quality of data extractions and resolved disagreements through discussion with the full review team (CS, BB, EP, MP and EA). Two senior reviewers in MH (MP and EA) supervised the extraction, randomly checking 20% of the extracted data. Given the nature of the study, ethics approval was not applicable. CS, BB and EP independently evaluated the quality of the included studies using the Newcastle-Ottawa Scale (NOS) for observational studies.³⁸ First, the full review team minimally adapted the NOS to the specificity of the research questions, and to the design and features of relevant primary studies. Next, a senior reviewer (EA) provided structured training and supervision to guide the screeners in applying the NOS appropriately, consistently and uniformly. Finally, the full review team conducted follow-up meetings to discuss uncertainties and resolve conflicts on the interpretation and adaptation of the scale.

Main outcomes and measures

The primary outcome of this review was the use of health services during the COVID-19 pandemic. We measured access as a single contact with a health service. The incidence rate (IR) of the use of health services was measured as the number of accesses (eg, number of admissions or treatments) registered in a specific time divided by contact months, that is, the product between the population size of the catchment area specific to each service and the number of months observed. The rate was quantified at a single time point to analyse both measures of services used in the pre-pandemic period (up until March 2020) and during the pandemic period. The pandemic period was counted as defined by the authors of the primary studies, ensuring it was close to the dates on which the WHO declared the outbreak.³⁹ We calculated the number of accesses, allowing for multiple accesses per person (recurrent events), while controlling for differences in the observation days. We calculated contacts day using the number of contacts with services and catchment areas in different periods. The secondary outcome was the change in the use of health services between the pre-pandemic and pandemic periods. This outcome was expressed as the IR ratio (IRR) and quantified as the number of accesses per time unit. We conducted two sets of meta-analyses. We used the Stata metan module⁴⁰ for statistical analyses. We used random effects DerSimonian-Laird (DL) inverse-variance models as we expected between-study heterogeneity due to the variety of populations, their characteristics and outcome measures. Heterogeneity across studies was calculated with statistics I^2 ⁴¹ and τ^2 and its statistical significance was assessed through Cochran's Q. For the primary outcome,

we pooled aggregated data on the use of health services; since we assumed a log-normal distribution, rates were transformed into logarithms. We computed the pooled IR from primary studies and calculated the 95% CI by choosing as a unit of measurement the contact-month rate for 10 000 inhabitants. For the secondary outcome, we pooled aggregated data on the during-pre ratio of rates of use of health services. We computed the pooled IRR of service used in the two periods with a related 95% CI and tested the hypothesis of no variation (IRR=1). To assess potential causes of heterogeneity, we conducted subgroup meta-analyses on the following variables: (a) population age groups, that is, children and adolescents (aged <18 years) versus adults (aged ≥18 years) versus mixed population age groups and (b) type of health service, that is, MH services versus general health services. We measured both within and between subgroups heterogeneity; the statistical significance of the latter was used to test for heterogeneous results across subgroups. As described in the protocol of this review, we collected data and performed a moderation analysis to evaluate whether the assessment of psychological distress or diagnosis of mental disorders during the pandemic moderated the use of health services. We will present such results elsewhere.

Patient and public involvement

Patients or the public were not involved in the design, conduct, reporting or dissemination plans of our research.

RESULTS

Characteristics of included studies

The electronic search yielded a total of 10 014 records. Before screening, we removed 5155 records because they were duplicates, leaving 4859 records for screening. We examined the full text of 580 studies and identified 281 as potentially eligible (figure 1). We included 136^{26 42–153} studies in the review, of which 44 were meta-analysed. References to selected excluded studies are reported in the online supplemental material. The online supplemental tables 1.1 and 1.2 display selected characteristics of included studies. Of the 136 included studies, 108 were cross-sectional, 27 were cohort and 1 was case-control. 121 studies were conducted in high-income countries, while 15 were conducted in low- and middle-income countries.³⁶ Nineteen studies recruited children and adolescents, 35 recruited adults and 82 recruited a mixed population in terms of age. Seventy-nine studies examined populations consisting of patients with pre-existing mental disorders (eg, CMDs, substance use, eating disorders), as well as those with other clinical conditions (eg, cancer, cardiovascular diseases, physical disabilities); 21 focused on the general population. Furthermore, 36 studies included both the general population and populations with pre-existing mental disorders or clinical conditions. Sixty-three studies were conducted in hospital settings and 28 in community health services. Forty-five were categorised as 'multicentre' because they were conducted in both




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 flow diagram for new systematic reviews, which included searches of databases ICTRP, International Clinical Trials Registry Platform; mental health.

hospital and community health settings. The outcome measures of 95 studies assessed the use of MH services, while 22 focused on the use of general health services. Nineteen examined the use of both general health and MH services. Of the studies included in the meta-analysis, mean observation days in the prepandemic period were 287 (range: 18–1461) and 227 (range: 28–490) in the pandemic period. The mean number of contacts in the prepandemic period was 661 649 (range: 25–27 224 568) and 582 442 (range 10–25 274 298) in the pandemic period. Studies did not vary substantially in quality (online supplemental tables 2–4): 28 studies met the criteria for quality assessment and 16 did not meet all quality criteria, still scoring a good overall.¹⁵⁴ The Egger’s test provided support to the unlikelihood of publication bias ($p=0.769$).

Rate of general health and MH services use

The meta-analysis of the primary outcome (44 studies) revealed that the IR of use of both general health and MH services during the COVID-19 pandemic was 2.59 contacts months per 10 000 inhabitants (overall IR=2.59; 95% CI: 1.30 to 5.16; $I^2=100%$) (figure 2). When exploiting Cochran’s Q statistic, we observed a statistically significant difference in effect sizes across all studies included in the overall meta-analysis ($\tau^2=5.44$; $p<0.001$).

Change in use of general health and MH services

When observing IRR between the prepandemic and the pandemic periods, the overall meta-analysis (42 studies) revealed a statistically significant ($p<0.001$) change in the use of both general health and MH services (overall




Figure 2 Rate of health services use (incidence rate). Overall meta-analysis pooling of aggregate data using the random-effects inverse-variance model with the DerSimonian-Laird (DL) estimate of τ^2 . Results are shown on a logarithmic scale.

IRR=0.75; 95% CI: 0.65 to 0.79; $I^2=99.9\%$) (figure 3). Thus, we identified a reduction of 28.5% in the use of health services during the pandemic. When exploiting Cochran's Q statistic, we observed a statistically significant difference in effect sizes across all included studies ($\tau^2=0.09$; $p<0.001$). The comparative magnitude of between-study heterogeneity was considerable.

Subgroup analysis

Primary outcome

Subgroup analyses are reported in the online supplemental material. The subgroup meta-analysis on the variable (a) revealed that the rate of the use of health services during the pandemic in studies including a mixed population in terms of age (IR=3.50; 95% CI: 1.51 to 8.14) was higher than the rate of studies involving children and adolescents (IR=1.02; 95% CI: 0.23 to 4.51) and higher than the rate of studies involving adults (IR=2.69; 95% CI: 0.90 to 8.04). When exploiting Cochran's Q statistic, we observed statistically significant differences in effect sizes within subgroups. Yet the between-subgroup heterogeneity calculated using DL subgroup weights was not statistically significant ($Q=2.01$; $df=2$; $p=0.366$) (online supplemental figure 1.1). The subgroup meta-analysis on the variable 'type of health services revealed that, during the pandemic, the rate of MH services use was significantly higher (IR=2.86; 95% CI: 1.42 to 5.77) than the general health services rate (IR=0.69; 95% CI: 0.31 to 1.55); here, the between-subgroup heterogeneity calculated using DL subgroup weights was statistically significant ($Q=6.76$; $df=1$; $p=0.009$). When exploiting Cochran's Q statistic, we observed statistically significant differences in effect sizes within the subgroups (online supplemental figure 1.2).

Secondary outcome

The subgroup meta-analysis on the variable (a) revealed that studies including adults reported a decrease in the use of health services in the pandemic period compared with the pre-pandemic period (IRR=0.80; 95% CI: 0.66 to 0.98). The decrease in use was lower when compared with the change observed in studies involving children and adolescents (IRR=0.73; 95% CI: 0.48 to 1.11) and the change observed in studies involving a mixed population in terms of age (IRR=0.68; 95% CI: 0.61 to 0.76). Yet, the estimate of studies comprising children and adolescents was not significant ($p=0.140$), while it was for adults ($p=0.027$) and mixed ($p<0.001$) subgroups. When exploiting Cochran's Q statistic, we observed statistically significant differences in effect sizes within subgroups. The between-subgroup heterogeneity calculated using DL subgroup weights was not statistically significant ($Q=2.09$; $df=2$; $p=0.35$) (online supplemental figure 2.1). The subgroup meta-analysis on the variable (b) revealed that MH services presented a statistically significant lower change in use between the pre-pandemic and pandemic periods (IRR=0.71; 95% CI: 0.65 to 0.78) when compared with the change observed in the general health services use (IRR=0.82; 95% CI: 0.53 to 1.25). The between-subgroup heterogeneity calculated

using DL subgroup weights was not statistically significant ($Q=0.38$; $df=1$; $p=0.53$). When testing subgroup effect size, there were statistically significant differences in effect sizes among the MH services subgroup ($p<0.001$), but no statistically significant differences in the effect size among the general health services subgroup ($p=0.35$). When exploiting Cochran's Q statistic, we observed statistically significant differences in effect sizes within the subgroups (online supplemental figure 2.2).

DISCUSSION

Our study provides a systematic synthesis of the evidence on the impact of the COVID-19 pandemic on the use of health services while accounting for MH. The 44 studies included in our meta-analysis were greatly heterogeneous and of adequate quality. The overall estimate revealed a reduction of 28.5% in the use of health services between the pre-pandemic and pandemic periods, with negligible and little difference by age group and type of service. Variance was fully dominated by its between-study component, also due to the population-level nature of the estimates, whose high precision led to a very low within-study variance. When meta-analysed subgroups did not meet statistical significance, such results are ascribable to the limited statistical power resulting from the small number of studies available within subgroups. The COVID-19 PH responses led to disruptions and reorganisations that affected the offer modalities, accessibility and availability of health services,²⁻⁵ and our study stands in as a sound demonstration of a reduction in the use of services during that time. Our results significantly expand current evidence on decreasing hospital admissions, emergency presentations and outpatient care.^{155 156} The drop in the use of services was ascertained both for physical health^{7 18} and for MH conditions.^{18 19 157 158} Reduced care contacts were reported in studies carried out in Europe and in other parts of the world^{4 12 13 159-161}; French studies showed declined admissions¹³ and halved emergency presentations for psychiatric disorders.¹⁶² Similar findings were obtained in two German studies reporting a 27% decrease in emergency cases¹⁶³ and a 25% reduction in hospitalisations following the introduction of the PH measures.¹⁶⁴ A study from the UK¹⁶ and a study from South Korea reported substantial reductions in outpatient care visits for MH conditions.¹⁷ Similar findings were obtained in a study from Canada revealing a decline in hospital admissions and emergency presentations in children and adolescents.¹⁶⁵ Two studies from Italy and one study from the UK unveiled curtailed surgical and diagnostic procedures¹⁶⁶⁻¹⁶⁸ across different care specialties. When looking at the standard framework for access to healthcare, which includes approachability and availability, as well as the ability to perceive and seek care,¹




Figure 3 Change in use of health services (incidence rate ratio). Overall meta-analysis pooling of aggregate data using the random-effects inverse-variance model with the DerSimonian-Laird (DL) estimate of τ^2 .



we ponder that: first, to reduce the spread of the virus, governments worldwide recommended citizens seek hospital care only when strictly necessary (ie, approachability)¹²; second, the fear of infection^{169 170} might have prevented patients from seeking and using care³³ (ie, ability to perceive and seek care) and third, the redirection of resources toward COVID-19 therapy resulted in de-prioritisation of routine care and elective interventions, lack of outpatient care and non-availability of treatment in hospital settings¹⁷⁰; this shift also led to restrictions or denials of access to healthcare (ie, accessibility).¹⁷¹ Reductions in access to and provision of services imply substantial diagnostic, treatment and care gaps, resulting in short-, mid- and long-term consequences for population health. These include increased morbidity and mortality, poorer clinical outcomes, delayed recovery and higher rates of medical relapse across various care specialties.^{18 171} Our results are important because they provide an accurate quantification of the change in the use of services that may be of relevance to model its impact on population health outcomes. By stratifying the use of general health and MH services, our results emphasise the need for attention to MH services that, though severely affected during the pandemic, still lack priority on the policy agenda.¹⁷² By focusing on a population with assessments of psychological distress or diagnosis of mental disorders, our review provides novel evidence for the direction of the association between MH and the use of health services. Unveiling this association is of particular importance because individuals with CMDs suffered disproportionately from the pandemic responses compared with the general population.¹⁶⁶ Discontinuation of MH services plausibly contributed to both the increase in the severity of CMDs and a greater impact of CMDs on socioeconomic and life circumstances, thus exacerbating pre-existing health inequalities in such a vulnerable population group.¹⁶⁶ An additional novelty of our review is that we retrieved studies from high and low- and middle-income countries, which confirmed that evidence is sparser in the latter than in the former setting.¹⁸ This is the first research to patch the void related to the unclear association between MH and the use of health services during the pandemic by focusing on a population with assessments of psychological distress or diagnosis of mental disorders and differentiating services. The internal validity of our study is corroborated by the advanced statistical analyses and the standardisation of procedures, which limited information bias, increased reliability and assured quality of data extraction and formal assessment of quality of studies. The generalisability across various services, countries, population groups and community characteristics of our findings relies on the extensive search strategy in databases, proactive and systematic contact with authors and the screening and selection steps done in

double. Nonetheless, several limitations should be acknowledged. The breadth of our review resulted in substantial heterogeneity in population data metrics and outcome measures across studies (eg, access measured at the patient level versus at the service level). Additionally, the size of catchment areas varied greatly across settings. Furthermore, the included studies were conducted during different pandemic waves and in various countries, which added complexity to the consistency of outcomes. Differences in virus spread, infection rates and PH restrictions across countries and regions further contributed to this variability. The heterogeneity of the collected data also precluded subgroup meta-analysis based on clinical characteristics of the population, such as pre-existing MH disorders or COVID-19 infection status. Another limitation of our review is that we restricted our search to five databases and applied language restrictions, which may have limited the comprehensiveness of our findings. Moreover, the generalisability of our results to patients without assessments of psychological distress or diagnosis of mental disorders is constrained by the specific focus we adopted. Due to the observational nature of our review, the interpretability of Egger's test is limited. Similarly, the interpretability of NOS score is limited because we adapted and standardised its use to align with the inclusion criteria of our meta-analysis, rather than the systematic review, and focused on how outcomes were measured in the primary studies. Finally, it should be noted that our search was completed in March 2023, and newer, potentially relevant studies may have since been published. To ensure the continued relevance of our findings, an update of this review and meta-analysis is planned for the end of 2025. The COVID-19 pandemic highlighted the necessity for PH responses to meet new population care needs while also addressing long-standing issues of scarce, unequal and inefficient healthcare resources.¹⁷³ The pandemic could provide a window of opportunity for promoting the reorganisation and reorientation of services towards needs-centred models. One example is the introduction of telemedicine, in the form of videoconferencing or phone, to permit follow-up of the patient. Telemedicine has set itself up as a practical and distinctive alternative to on-site care.^{12 57} Previous studies have demonstrated its efficacy in managing chronic and MH care, suggesting the potential for long-lasting implementation. Still, the implications of its use on the quantity and quality of services supplied must be thoroughly explored.

CONCLUSION

While considering MH, COVID-19 has led to a decline in the use of health services, with minimal variation across age groups and service types. Significant

reductions in access to and provision of services led to unmet care needs, resulting in unpredictable consequences for population health outcomes.

Acknowledgements The authors of this manuscript extend their sincere gratitude to all corresponding authors who responded to their outreach and generously provided them with the necessary data to perform the meta-analysis. Namely: Giulio Castelpietra, Marcos Del Pozo Banos, Harriet Hiscock, Levy Jäger, Ryan Loh, Christel M. Middeldorp, Aleksandar Savic, Hannah Shaw, Scott A. Simpson, Hande Ayraler Taner, Jakša Vukojević and Jane Zhu.

Contributors CS and EA conceptualised the study. MBB provided support with the conceptualisation of the study and the definition of the research questions. CS defined the search strategy. CS, BB and EP performed the study selection, data extraction and data curation. CB and MP conceptualised the methodology and supervised the study selection and data extraction process. FT, CB and MP performed the formal statistical analysis. FT did the visualisation of the meta-analysis outputs. CS, BB and EP assisted throughout the statistical analysis. CS wrote the original draft. MP and EA reviewed and edited the manuscript. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication. CS and EA directly accessed and verified the underlying data reported in the manuscript. CS is responsible for the overall content as guarantor.

Funding This research was supported by the Università della Svizzera Italiana (USI) and the State Secretariat of Education, Research and Innovation (SERI) in Switzerland through its project-related contribution P-4 Swiss Learning Health System (SLHS). See also: <https://www.slhs.ch>

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting, dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Data collected for this study, including the full list of included and excluded records, raw extracted data and meta-analysed data will be made available on request by contacting the corresponding author. The review protocol, search string used to retrieve data, list of selected excluded studies, tables 1.1 and 1.2, and subgroup meta-analysis are available in the online supplemental material.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Camilla Sculco <http://orcid.org/0009-0005-0735-9360>

Eleonora Prina <http://orcid.org/0000-0002-4005-1967>

Marianna Purgato <http://orcid.org/0000-0002-3783-8195>

REFERENCES

- Cu A, Meister S, Lefebvre B, *et al*. Assessing healthcare access using the Levesque's conceptual framework- a scoping review. *Int J Equity Health* 2021;20:116.
- WHO. Mental health and covid-19: early evidence of the pandemic's impact. 2022.
- Thome J, Deloyer J, Coogan AN, *et al*. The impact of the early phase of the COVID-19 pandemic on mental-health services in Europe. *World J Biol Psychiatry* 2021;22:516–25.
- Yalçın M, Baş A, Bilici R, *et al*. Psychiatric emergency visit trends and characteristics in a mental health epicenter in Istanbul during COVID-19 lockdown. *Soc Psychiatry Psychiatr Epidemiol* 2021;56:2299–310.
- Bakolis I, Stewart R, Baldwin D, *et al*. Changes in daily mental health service use and mortality at the commencement and lifting of COVID-19 'lockdown' policy in 10 UK sites: a regression discontinuity in time design. *BMJ Open* 2021;11:e049721.
- Deml MJ, Minnema J, Dubois J, *et al*. The impact of the COVID-19 pandemic on the continuity of care for at-risk patients in Swiss primary care settings: A mixed-methods study. *Soc Sci Med* 2022;298:114858.
- Jeffery MM, D'Onofrio G, Paek H, *et al*. Trends in Emergency Department Visits and Hospital Admissions in Health Care Systems in 5 States in the First Months of the COVID-19 Pandemic in the US. *JAMA Intern Med* 2020;180:1328–33.
- Esenwa C, Parides MK, Labovitz DL. The effect of COVID-19 on stroke hospitalizations in New York City. *J Stroke Cerebrovasc Dis* 2020;29:105114.
- Pelletier JH, Rakkar J, Au AK, *et al*. Trends in US Pediatric Hospital Admissions in 2020 Compared With the Decade Before the COVID-19 Pandemic. *JAMA Netw Open* 2021;4:e2037227.
- Boldrini T, Girardi P, Clerici M, *et al*. Consequences of the COVID-19 pandemic on admissions to general hospital psychiatric wards in Italy: Reduced psychiatric hospitalizations and increased suicidality. *Prog Neuropsychopharmacol Biol Psychiatry* 2021;110:110304.
- Wyatt S, Mohammed MA, Fisher E, *et al*. Impact of the SARS-CoV-2 pandemic and associated lockdown measures on attendances at emergency departments in English hospitals: A retrospective database study. *Lancet Reg Health Eur Mar* 2021;2:100034.
- Gómez-Ramiro M, Fico G, Anmella G, *et al*. Changing trends in psychiatric emergency service admissions during the COVID-19 outbreak: Report from a worldwide epicentre. *J Affect Disord* 2021;282:26–32.
- Pignon B, Gourevitch R, Tebeka S, *et al*. Dramatic reduction of psychiatric emergency consultations during lockdown linked to COVID-19 in Paris and suburbs. *Psychiatry Clin Neurosci Oct* 2020;74:557–9.
- Balestrieri M, Rucci P, Amendola D, *et al*. Emergency Psychiatric Consultations During and After the COVID-19 Lockdown in Italy. A Multicentre Study. *Front Psychiatry* 2021;12:697058.
- McDowell MJ, Fry CE, Nisavic M, *et al*. Evaluating the association between COVID-19 and psychiatric presentations, suicidal ideation in an emergency department. *PLoS One* 2021;16:e0253805.
- Mansfield KE, Mathur R, Tazare J, *et al*. Indirect acute effects of the COVID-19 pandemic on physical and mental health in the UK: a population-based study. *Lancet Digit Health* 2021;3:e217–30.
- Seo JH, Kim SJ, Lee M, *et al*. Impact of the COVID-19 pandemic on mental health service use among psychiatric outpatients in a tertiary hospital. *J Affect Disord* 2021;290:279–83.
- Wettstein A, Tlali M, Joska JA, *et al*. The effect of the COVID-19 lockdown on mental health care use in South Africa: an interrupted time-series analysis. *Epidemiol Psychiatr Sci* 2022;31:e43.
- Williams R, Jenkins DA, Ashcroft DM, *et al*. Diagnosis of physical and mental health conditions in primary care during the COVID-19 pandemic: a retrospective cohort study. *Lancet Public Health* 2020;5:e543–50.
- Aragona M, Barbato A, Cavani A, *et al*. Negative impacts of COVID-19 lockdown on mental health service access and follow-up adherence for immigrants and individuals in socio-economic difficulties. *Public Health (Fairfax)* 2020;186:52–6.
- Ougrin D, Wong BH, Vaezinejad M, *et al*. Pandemic-related emergency psychiatric presentations for self-harm of children and adolescents in 10 countries (PREP-kids): a retrospective international cohort study. *Eur Child Adolesc Psychiatry* 2022;31:1–13.
- Fiorillo A, Sampogna G, Giallonardo V, *et al*. Effects of the lockdown on the mental health of the general population during the COVID-19 pandemic in Italy: Results from the COMET collaborative network. *Eur Psychiatr* 2020;63:e87.
- Alleaume C, Verger P, Peretti-Watel P, *et al*. Psychological support in general population during the COVID-19 lockdown in France: Needs and access. *PLoS One* 2021;16:e0251707.
- Brooks SK, Webster RK, Smith LE, *et al*. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020;395:912–20.
- Winkler P, Formanek T, Mlada K, *et al*. Increase in prevalence of current mental disorders in the context of COVID-19: analysis of



- repeated nationwide cross-sectional surveys. *Epidemiol Psychiatr Sci* 2020;29:e173.
- 26 Bu F, Mak HW, Fancourt D. Rates and predictors of uptake of mental health support during the COVID-19 pandemic: an analysis of 26,720 adults in the UK in lockdown. *Soc Psychiatry Psychiatr Epidemiol* 2021;56:2287–97.
 - 27 Liao YH, Fan BF, Zhang HM, *et al.* The impact of COVID-19 on subthreshold depressive symptoms: a longitudinal study. *Epidemiol Psychiatr Sci* 2021;30:e20.
 - 28 Chow MW, Noorthoorn EO, Wierdsma AI, *et al.* Impact of the first COVID-19 outbreak on mental health service utilisation at a Dutch mental health centre: retrospective observational study. *BJPsych Open* 2021;7:e213.
 - 29 Moreno C, Wykes T, Galderisi S, *et al.* How mental health care should change as a consequence of the COVID-19 pandemic. *Lancet Psychiatry* 2020;7:813–24.
 - 30 Benuto LT, Casas J, Gonzalez F, *et al.* The Behavioral Model of Health: Education, Behavioral Health Factors, and Stigma as Predictors of Help-Seeking Attitudes. *Community Ment Health J* 2020;56:1275–83.
 - 31 Richter D, Bonsack C, Burr C, *et al.* Therapeutic alliance, social inclusion and infection control—Towards pandemic-adapted mental healthcare services in Switzerland. Swiss Archives of Neurology, Psychiatry and Psychotherapy. *Int J Health Policy Manag* 2021;12.
 - 32 Rachamin Y, Meyer MR, Rosemann T, *et al.* Impact of the COVID-19 Pandemic on Elective and Emergency Inpatient Procedure Volumes in Switzerland - A Retrospective Study Based on Insurance Claims Data. *Int J Health Policy Manag* 2023;12:6932.
 - 33 Rezapour R, Dorosti AA, Farahbakhsh M, *et al.* The impact of the Covid-19 pandemic on primary health care utilization: an experience from Iran. *BMC Health Serv Res* 2022;22:404.
 - 34 Association AP. *Diagnostic and statistical manual of mental disorders*. 5th edn. 2022.
 - 35 The World Bank. World bank country and lending groups. Available: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> [Accessed 1 Sep 2023].
 - 36 EndNote. *Version EndNote 20*. Clarivate, 2013.
 - 37 Rayyan. Rayyan — a web and mobile app for systematic reviews. 2016.
 - 38 Wells GA, Wells G, Shea B, *et al.* The newcastle-ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2014.
 - 39 Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Biomed* 2020;91:157–60.
 - 40 METAN. *Stata module for fixed and random effects meta-analysis*. 2006.
 - 41 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
 - 42 Dragovic M, Pascu V, Hall T, *et al.* Emergency department mental health presentations before and during the COVID-19 outbreak in Western Australia. *Australas Psychiatry* 2020;28:627–31.
 - 43 Abbas MJ, Kronenberg G, McBride M, *et al.* The Early Impact of the COVID-19 Pandemic on Acute Care Mental Health Services. *Psychiatr Serv* 2021;72:242–6.
 - 44 Adams SH, Schaub JP, Nagata JM, *et al.* Young Adult Anxiety or Depressive Symptoms and Mental Health Service Utilization During the COVID-19 Pandemic. *J Adolesc Health* 2022;70:985–8.
 - 45 Alsaïdi I, De Sousa Santos F, Plard B, *et al.* Factors associated with SARS-CoV2 infection and care pathways among the most vulnerable populations living in Marseille: a case control study. *BMC Public Health* 2021;21:1704.
 - 46 Amram O, Amiri S, Panwala V, *et al.* The impact of relaxation of methadone take-home protocols on treatment outcomes in the COVID-19 era. *Am J Drug Alcohol Abuse* 2021;47:722–9.
 - 47 Ayraler Taner H, Akin Sari B, Kaba D. How did the COVID-19 pandemic affect child and adolescent psychiatry outpatient clinic admissions? A single-center, retrospective study. *Clin Child Psychol Psychiatry* 2022;27:824–35.
 - 48 Ayton A, Viljoen D, Ryan S, *et al.* Risk, demand, capacity and outcomes in adult specialist eating disorder services in South-East of England before and since COVID-19. *BJPsych Bull* 2022;46:89–95.
 - 49 Banks DE, Paschke ME, Li X, *et al.* Opioid Use Disorder and COVID-19: Treatment and Recovery Factors among Vulnerable Populations at the Intersection of Two U.S. Epidemics. *J Psychoactive Drugs* 2022;54:300–8.
 - 50 Barlati S, Nibbio G, Bianchi F, *et al.* Long-acting Injectable Antipsychotics during the COVID-19 pandemic in schizophrenia: An observational study in a real-world clinical setting. *Psychiatry Res* 2022;317:114878.
 - 51 Barnett ML, Waken RJ, Zheng J, *et al.* Changes in Health and Quality of Life in US Skilled Nursing Facilities by COVID-19 Exposure Status in 2020. *JAMA* 2022;328:941–50.
 - 52 Barrett EA, Simonsen C, Aminoff SR, *et al.* The COVID-19 pandemic impact on wellbeing and mental health in people with psychotic and bipolar disorders. *Brain Behav* 2022;12:e2559.
 - 53 Bauer-Staeb C, Davis A, Smith T, *et al.* The early impact of COVID-19 on primary care psychological therapy services: A descriptive time series of electronic healthcare records. *EClinicalMedicine* 2021;37:100939.
 - 54 Beghi M, Ferrari S, Biondi L, *et al.* Mid-term psychiatric consequences of the COVID-19 pandemic: a 4 months observational study on emergency room admissions for psychiatric evaluation after the (first) lockdown period in Italy. *Soc Psychiatry Psychiatr Epidemiol* 2022;57:1283–9.
 - 55 Bohlken J, Kostev K, Riedel-Heller S, *et al.* Effect of the COVID-19 pandemic on stress, anxiety, and depressive disorders in German primary care: A cross-sectional study. *J Psychiatr Res* 2021;143:43–9.
 - 56 Bothara RK, Raina A, Carne B, *et al.* Paediatric presentations to Christchurch Hospital Emergency Department during COVID-19 lockdown. *J Paediatr Child Health* 2021;57:877–82.
 - 57 Brandizzi M, PolSELLI A, Corigliano V, *et al.* Psychiatric emergencies during, after, and before the COVID-19 lockdown: what happened to our patients? A naturalistic observational study. *Ann Gen Psychiatry* 2022;21:29.
 - 58 Brausch AM, Whitfield M, Clapham RB. Comparisons of mental health symptoms, treatment access, and self-harm behaviors in rural adolescents before and during the COVID-19 pandemic. *Eur Child Adolesc Psychiatry* 2023;32:1051–60.
 - 59 Brothers S, Viera A, Heimer R. Changes in methadone program practices and fatal methadone overdose rates in Connecticut during COVID-19. *J Subst Abuse Treat* 2021;131:108449.
 - 60 Bruckner TA, Huo S, Huynh M, *et al.* Psychiatric Emergencies in Los Angeles County During, and After, Initial COVID-19 Societal Restrictions: An Interrupted Time-Series Analysis. *Community Ment Health J* 2023;59:622–30.
 - 61 Cameron EE, Joyce KM, Delaquis CP, *et al.* Maternal psychological distress & mental health service use during the COVID-19 pandemic. *J Affect Disord* 2020;276:765–74.
 - 62 Cann J, Barter R, Battle J, *et al.* Patients presenting to an acute general hospital with acute mental health needs: a retrospective observational cohort study. *BMJ Open* 2022;12:e059102.
 - 63 Carr MJ, Steeg S, Webb RT, *et al.* Effects of the COVID-19 pandemic on primary care-recorded mental illness and self-harm episodes in the UK: a population-based cohort study. *Lancet Public Health* 2021;6:e124–35.
 - 64 Castelpietra G, Colli C, Tossut D, *et al.* The impact of Covid-19 pandemic on community-oriented mental health services: The experience of Friuli Venezia Giulia region, Italy. *Health Policy Technol* 2021;10:143–50.
 - 65 Chakraborty N, Carr R, Tromans S. The impact of covid -19 on an early intervention in psychosis service . *Prog Neurol Psychiatry* 2020;24:17–21.
 - 66 Chakrapani V, Newman PA, Sebastian A, *et al.* Mental health, economic well-being and health care access amid the COVID-19 pandemic: a mixed methods study among urban men who have sex with men in India. *Sexual and Reproductive Health Matters* 2022;30:2144087.
 - 67 Choi SH, Kim S. Factors associated with adverse mental health outcomes during the COVID-19 pandemic. *Public Health Pract (Oxf)* 2023;5:100360.
 - 68 Cohen KA, Manikandan D, Jirsa M, *et al.* Mental healthcare on college campuses during COVID-19: Comparing telehealth, in-person, and hybrid modes of delivery. *J Am Coll Health* 2024;72:3313–21.
 - 69 DelPozo-Banos M, Lee SC, Friedmann Y, *et al.* Healthcare contacts with self-harm during COVID-19: An e-cohort whole-population-based study using individual-level linked routine electronic health records in Wales, UK, 2016–March 2021. *PLoS ONE* 2016;17:e0266967.
 - 70 Di Lorenzo R, Pinelli M, Bertani D, *et al.* The Impact of COVID-19 Pandemic on Psychiatric Emergencies in Two Different Settings: Emergency Room and Community Mental Health Service. *Front Psychiatry* 2022;13:894939.
 - 71 Ekinci O, Erkan Ekinci A. Demographic and Clinical Factors Related to Severe COVID-19 Infection and Mortality in Patients With Schizophrenia. *J Nerv Ment Dis* 2022;210:257–63.
 - 72 Evensen M, Hart RK, Godøy AA, *et al.* Impact of the COVID-19 pandemic on mental healthcare consultations among children and adolescents in Norway: a nationwide registry study. *Eur Child Adolesc Psychiatry* 2023;32:1025–35.

- 73 Favreau M, Hillert A, Osen B, *et al.* Psychological consequences and differential impact of the COVID-19 pandemic in patients with mental disorders. *Psychiatry Res* 2021;302:114045.
- 74 Fond G, Pauly V, Leone M, *et al.* Disparities in Intensive Care Unit Admission and Mortality Among Patients With Schizophrenia and COVID-19: A National Cohort Study. *Schizophr Bull* 2021;47:624–34.
- 75 Geulayov G, Borschmann R, Mansfield KL, *et al.* Utilization and Acceptability of Formal and Informal Support for Adolescents Following Self-Harm Before and During the First COVID-19 Lockdown: Results From a Large-Scale English Schools Survey. *Front Psychiatry* 2022;13:881248.
- 76 Greig F, Perera G, Tsamakidis K, *et al.* Loneliness in older adult mental health services during the COVID-19 pandemic and before: Associations with disability, functioning and pharmacotherapy. *Int J Geriatr Psychiatry* 2021;37.
- 77 Grosland M, Larsen VB, Reneflot A, *et al.* The COVID-19 pandemic's impact on mental health care use among Norwegian students in higher education: a nation-wide register-based pre-post cohort study. *BMC Health Serv Res* 2022;22:1506.
- 78 Haderlein TP, Wong MS, Yuan A, *et al.* Association of PTSD with COVID-19 testing and infection in the Veterans Health Administration. *J Psychiatr Res* 2021;143:504–7.
- 79 Hedberg P, Granath F, Bruchfeld J, *et al.* Post COVID-19 condition diagnosis: A population-based cohort study of occurrence, associated factors, and healthcare use by severity of acute infection. *J Intern Med* 2023;293:246–58.
- 80 Henry N, Parthiban S, Farroha A. The effect of COVID-19 lockdown on the incidence of deliberate self-harm injuries presenting to the emergency room. *Int J Psychiatry Med* 2021;56:266–77.
- 81 Hernández-Calle D, Andreo-Jover J, Curto-Ramos J, *et al.* Pediatric Mental Health Emergency Visits During the COVID-19 Pandemic. *Scand J Child Adolesc Psychiatr Psychol* 2022;10:53–7.
- 82 Hildersley R, Easter A, Bakolis I, *et al.* Changes in the identification and management of mental health and domestic abuse among pregnant women during the COVID-19 lockdown: regression discontinuity study. *BJPsych open* 2022;8:e96.
- 83 Hirata R, Kawashima H, Tsuboi T, *et al.* An Online Survey About Electroconvulsive Therapy in Japan During the COVID-19 Pandemic: Comparison of Early and Recent Stages. *Neuropsychiatr Dis Treat* 2022;18:1277–85.
- 84 Hiscock H, Chu W, O'Reilly G, *et al.* Association between COVID-19 restrictions and emergency department presentations for paediatric mental health in Victoria, Australia. *Aust Health Rev* 2022;46:529–36.
- 85 Hoffman KA, Foot C, Levander XA, *et al.* Treatment retention, return to use, and recovery support following COVID-19 relaxation of methadone take-home dosing in two rural opioid treatment programs: A mixed methods analysis. *J Subst Abuse Treat* 2022;141:108801.
- 86 Holland KM, Jones C, Vivolo-Kantor AM, *et al.* Trends in US Emergency Department Visits for Mental Health, Overdose, and Violence Outcomes Before and During the COVID-19 Pandemic. *JAMA Psychiatry* 2021;78:372–9.
- 87 Hooper K, Hooper M, Nguyen J, *et al.* Are you vaccinated? COVID-19 vaccination rates and the effect of a vaccination program in a metropolitan mental health inpatient population in Australia. *Australas Psychiatry* 2023;31:38–42.
- 88 Joo SW, Kim H, Jo YT, *et al.* Impact of the early phase of the COVID-19 pandemic on the use of mental health services in South Korea: a nationwide, health insurance data-based study. *Soc Psychiatry Psychiatr Epidemiol* 2023;58:441–52.
- 89 Kadra-Scalzo G, Kornblum D, Stewart R, *et al.* Adverse outcomes associated with recorded victimization in mental health electronic records during the first UK COVID-19 lockdown. *Soc Psychiatry Psychiatr Epidemiol* 2023;58:431–40.
- 90 Kavanagh A, Hatton C, Stancliffe RJ, *et al.* Health and healthcare for people with disabilities in the UK during the COVID-19 pandemic. *Disabil Health J* 2022;15:101171.
- 91 Kendzerska T, Zhu DT, Pugliese M, *et al.* Trends in all-cause mortality and inpatient and outpatient visits for ambulatory care sensitive conditions during the first year of the COVID-19 pandemic: A population-based study. *J Hosp Med* 2022;17:726–37.
- 92 Köse S, Özbaran B, Doğan N, *et al.* Are the illness severity and treatment efficacy of the patients followed in the child psychiatry inpatient service during the pandemic period different from the Normal period? *J Clin Psy* 2022;25:168–76.
- 93 Krieger I, Erez G, Weinstein O, *et al.* COVID-19 Morbidity Among Individuals with Autistic Spectrum Disorder: A Matched Controlled Population-Based Study. *J Autism Dev Disord* 2023;53:789–94.
- 94 Lantos JD, Yeh HW, Raza F, *et al.* Suicide Risk in Adolescents During the COVID-19 Pandemic. *Pediatrics* 2022;149:2021–053486.
- 95 Leff RA, Setzer E, Cicero MX, *et al.* Changes in pediatric emergency department visits for mental health during the COVID-19 pandemic: A cross-sectional study. *Clin Child Psychol Psychiatry* 2021;26:33–8.
- 96 Lega I, Bramante A, Lauria L, *et al.* The Psychological Impact of COVID-19 among Women Accessing Family Care Centers during Pregnancy and the Postnatal Period in Italy. *Int J Environ Res Public Health* 2022;19:1983.
- 97 Lešin Gačina D, Jandroković S, Marčinko D, *et al.* Anxiety and Treatment Adherence among Glaucoma Patients during COVID-19 Pandemic and Earthquakes in Croatia. *Psychiatr Danub* 2022;34:348–55.
- 98 Li H, Chen X, Pan J, *et al.* Impact of the COVID-19 Pandemic on Utilization of Inpatient Mental Health Services in Shanghai, China. *Healthcare (Basel)* 2022;10:1402.
- 99 Liberman JN, Bhattacharjee S, Rui P, *et al.* Impact of the COVID-19 Pandemic on Healthcare Resource Utilization in Individuals with Major Depressive Disorder. *Health Serv Res Manag Epidemiol* 2022;9:2333392822111864.
- 100 Liu C-H, Chen P-C, Chen J-H, *et al.* Changes in self-harm- and violence-related urgent psychiatric consultation in the emergency department during the different stages of the COVID-19 pandemic. *BMC Psychiatry* 2022;22:384.
- 101 Loeb TB, Vidulich I, Smith-Clapham AM, *et al.* Unmet need for mental health services utilization among under-resourced Black and Latinx adults. *Fam Syst Health* 2023;41:149–59.
- 102 Luca L, Ciubara AB, Fulga I, *et al.* Social Implications for Psychiatric Pathology of Depressive and Anxiety Disorders, Alcohol Addiction and Psychotic Disorders during the COVID-19 Pandemic in Romania. Analysis of two Relevant Psychiatry Hospitals. *RCIS* 2020;69:261–72.
- 103 Mancheño-Velasco C, Dacosta-Sánchez D, Blanc-Molina A, *et al.* Changes in the Care Activity in Addiction Centers with Dual Pathology Patients during the COVID-19 Pandemic. *J Clin Med* 2022;11:4341.
- 104 McGuire AB, Flanagan ME, Kukla M, *et al.* n.d. Inpatient Mental Healthcare before and during the COVID-19 Pandemic. *Healthcare (Basel)* 9:1613.
- 105 McNicholas F, Kelleher I, Hedderman E, *et al.* Referral patterns for specialist child and adolescent mental health services in the Republic of Ireland during the COVID-19 pandemic compared with 2019 and 2018. *BJPsych Open* 2021;7:e91.
- 106 Menculini G, Tortorella A, Albert U, *et al.* Access to Mental Health Care during the First Wave of the COVID-19 Pandemic in Italy: Results from the COMET Multicentric Study. *Brain Sci* 2021;11:1413.
- 107 Mevorach T, Zur G, Benaroya-Milshtein N, *et al.* A Following Wave Pattern of Suicide-Related Pediatric Emergency Room Admissions during the COVID-19 Pandemic. *Int J Environ Res Public Health* 2023;20:1619.
- 108 Montalbani B, Bargagna P, Mastrangelo M, *et al.* The COVID-19 Outbreak and Subjects With Mental Disorders Who Presented to an Italian Psychiatric Emergency Department. *J Nerv Ment Dis* 2021;209:246–50.
- 109 Muhajarine N, Adeyinka DA, Pisolkar V, *et al.* Equity Analysis of Repeated Cross-Sectional Survey Data on Mental Health Outcomes in Saskatchewan, Canada during COVID-19 Pandemic. *Int J Environ Res Public Health* 2022;19:13808.
- 110 Mukadam N, Sommerlad A, Wright J, *et al.* Acute mental health presentations before and during the COVID-19 pandemic. *BJPsych Open* 2021;7:e134.
- 111 Nantaayi B, Ndawula RK, Musoke P, *et al.* Psychological Distress and Access to Mental Health Services Among Undergraduate Students During the COVID-19 Lockdown in Uganda. *Front Psychiatry* 2022;13:792217.
- 112 Nejati N, Crocker C, Kolajova M, *et al.* Examination of the impact of COVID-19 public health quarantine measures on acute mental health care services: A retrospective observational study. *Psychiatry Res* 2021;302:113999.
- 113 Nguyen KH, Chen S, Morris K, *et al.* Mental health symptoms and association with COVID-19 vaccination receipt and intention to vaccinate among adults, United States. *Prev Med* 2022;154:106905.
- 114 Norwood J, Kheshti A, Shepherd BE, *et al.* The Impact of COVID-19 on the HIV Care Continuum in a Large Urban Southern Clinic. *AIDS Behav* 2022;26:2825–9.
- 115 Olié E, Nogue E, Picot MC, *et al.* Could the decrease of the number of hospitalizations for suicide attempts in France be related to limited access to care? *J Affect Disord* 2022;302:313–4.

- 116 Ornell F, Borelli WV, Benzano D, *et al.* The next pandemic: impact of COVID-19 in mental healthcare assistance in a nationwide epidemiological study. *Lancet Reg Health Am* 2021;4:100061.
- 117 Payne N, van den Akker O. Mental health and coping with fertility treatment cessation during the COVID-19 pandemic in the UK. *J Psychosom Obstet Gynaecol* 2022;43:550–6.
- 118 Penington E, Lennox B, Geulayov G, *et al.* The early impact of the COVID-19 pandemic on patients with severe mental illness: An interrupted time-series study in South-East England. *Eur Psychiatry* 2022;65:e31.
- 119 Pérez V, Elices M, Vilagut G, *et al.* Suicide-related thoughts and behavior and suicide death trends during the COVID-19 in the general population of Catalonia, Spain. *Eur Neuropsychopharmacol* 2022;56:4–12.
- 120 Picchio CA, Valencia J, Doran J, *et al.* The impact of the COVID-19 pandemic on harm reduction services in Spain. *Harm Reduct J* 2020;17:87.
- 121 Quintas J, Guerreiro A, de Carvalho MJL, *et al.* The Implication of the First Wave of COVID-19 on Mental Health: Results from a Portuguese Sample. *Int J Environ Res Public Health* 2022;19:6489.
- 122 Rachamin Y, Jäger L, Schweighoffer R, *et al.* The Impact of COVID-19 on Mental Healthcare Utilization in Switzerland Was Strongest Among Young Females-Retrospective Study in 2018–2020. *Int J Public Health* 2023;68:1605839.
- 123 Rahman F, Evans N, Bernhardt J. Access to OUD Treatment and Maintenance of Sobriety amid the COVID-19 Pandemic. *Substance Use & Misuse* 2021;56:1005–9.
- 124 Rauf B, Sheikh H, Majid H, *et al.* COVID-19-related prescribing challenge in intellectual disability. *BJPsych open* 2021;7:e66.
- 125 Reece L, Sams DP. The impact of COVID-19 on adolescent psychiatric inpatient admissions. *Clin Child Psychol Psychiatry* 2022;27:112–21.
- 126 Roberti E, Giacchero R, Grumi S, *et al.* Post-partum Women's Anxiety and Parenting Stress: Home-Visiting Protective Effect During the COVID-19 Pandemic. *Matern Child Health J* 2022;26:2308–17.
- 127 Robillard R, Daros AR, Phillips JL, *et al.* Emerging New Psychiatric Symptoms and the Worsening of Pre-existing Mental Disorders during the COVID-19 Pandemic: A Canadian Multisite Study: Nouveaux symptômes psychiatriques émergents et détérioration des troubles mentaux préexistants durant la pandémie de la COVID-19: une étude canadienne multisite. *Can J Psychiatry* 2021;66:815–26.
- 128 Rømer TB, Christensen RHB, Blomberg SN, *et al.* Psychiatric Admissions, Referrals, and Suicidal Behavior Before and During the COVID-19 Pandemic in Denmark: A Time-Trend Study. *Acta Psychiatr Scand* 2021;144:553–62.
- 129 Ruprecht MM, Wang X, Johnson AK, *et al.* Evidence of Social and Structural COVID-19 Disparities by Sexual Orientation, Gender Identity, and Race/Ethnicity in an Urban Environment. *J Urban Health* 2021;98:27–40.
- 130 Sakowicz A, Matovina CN, Imeroni SK, *et al.* The association between the COVID-19 pandemic and postpartum care provision. *Am J Obstet Gynecol MFM* 2021;3:100460.
- 131 Sanchez K, da Graca B, Hall LR, *et al.* The Pandemic Experience for People with Depressive Symptoms: Substance Use, Finances, Access to Treatment, and Trusted Sources of Information. *Subst Abuse* 2022;16:11782218221126973.
- 132 Schers H, van Weel C, van Boven K, *et al.* The COVID-19 Pandemic in Nijmegen, the Netherlands: Changes in Presented Health Problems and Demand for Primary Care. *Ann Fam Med* 2021;19:44–7.
- 133 Schlegl S, Maier J, Meule A, *et al.* Eating disorders in times of the COVID-19 pandemic-Results from an online survey of patients with anorexia nervosa. *Int J Eat Disord* 2020;53:1791–800.
- 134 Seifert J, Meissner C, Birkenstock A, *et al.* Peripandemic psychiatric emergencies: impact of the COVID-19 pandemic on patients according to diagnostic subgroup. *Eur Arch Psychiatry Clin Neurosci* 2021;271:259–70.
- 135 Sharma M, Kataria D, Raut N, *et al.* Impact of COVID-19 pandemic on treatment seeking in opioid users: A retrospective comparative study. *Annals of Indian Psychiatry* 2022;6.
- 136 Shaw H, Robertson S, Ranceva N. What was the impact of a global pandemic (COVID-19) lockdown period on experiences within an eating disorder service? A service evaluation of the views of patients, parents/carers and staff. *J Eat Disord* 2021;9:14.
- 137 Spigel R, Lin JA, Milliren CE, *et al.* Access to care and worsening eating disorder symptomatology in youth during the COVID-19 pandemic. *J Eat Disord* 2021;9:69.
- 138 Steeg S, Carr M, Trefan L, *et al.* Primary care clinical management following self-harm during the first wave of COVID-19 in the UK: population-based cohort study. *BMJ Open* 2022;12:e052613.
- 139 Tedja AM, Shanmugam MS, Stathis S, *et al.* Short research article: COVID-19 and its impact on child and youth mental health service demand in the community and emergency department. *Child Adolesc Ment Health* 2023;28:167–71.
- 140 Toulany A, Kurdyak P, Guttman A, *et al.* Acute Care Visits for Eating Disorders Among Children and Adolescents After the Onset of the COVID-19 Pandemic. *J Adolesc Health* 2022;70:42–7.
- 141 Tromans S, Chester V, Harrison H, *et al.* Patterns of use of secondary mental health services before and during COVID-19 lockdown: observational study. *BJPsych Open* 2020;6:e117.
- 142 Tzur Bitan D, Kridin K, Cohen AD, *et al.* COVID-19 hospitalisation, mortality, vaccination, and postvaccination trends among people with schizophrenia in Israel: a longitudinal cohort study. *Lancet Psychiatry* 2021;8:901–8.
- 143 Vukojević J, Đuran N, Žaja N, *et al.* 100 Years apart: Psychiatric admissions during Spanish flu and COVID-19 pandemic. *Psychiatry Res* 2021;303:114071.
- 144 Wang J, Spencer A, Hulme C, *et al.* Healthcare utilisation, physical activity and mental health during COVID-19 lockdown: an interrupted time-series analysis of older adults in England. *Eur J Ageing* 2022;19:1617–30.
- 145 Winter BL, Hovaguimian F, Kouyos RD, *et al.* Changes in mental and sexual health among MSM using HIV pre-exposure prophylaxis during the SARS-CoV-2 pandemic: longitudinal analysis of the SwissPrEPared cohort study. *Swiss Med Wkly* 2022;152:w30192.
- 146 Yang J, Landrum MB, Zhou L, *et al.* Disparities in outpatient visits for mental health and/or substance use disorders during the COVID surge and partial reopening in Massachusetts. *Gen Hosp Psychiatry* 2020;67:100–6.
- 147 Yazkan Akgül G, Yıldırım Budak B, Burcu Erdoğan A, *et al.* Child and Adolescent Psychiatry Outpatient Clinic Referrals During Covid-19 Pandemic in Turkey. *Psychiatry Clin Psychopharmacol* 2022;32:140–8.
- 148 Yeoh E, Tan SG, Lee YS, *et al.* Impact of COVID-19 and partial lockdown on access to care, self-management and psychological well-being among people with diabetes: A cross-sectional study. *Int J Clin Pract* 2021;75:e14319.
- 149 Zhong B-L, Zhou D-Y, He M-F, *et al.* Mental health problems, needs, and service use among people living within and outside Wuhan during the COVID-19 epidemic in China. *Ann Transl Med* 2020;8:1392.
- 150 Zhu JM, Myers R, McConnell KJ, *et al.* Trends In Outpatient Mental Health Services Use Before And During The COVID-19 Pandemic. *Health Aff (Millwood)* 2022;41:573–80.
- 151 OD D, OD M, L D. Covid-19 and the Impact on Referrals to Psychiatry in those 65 years and Older. *Ir Med J* 2022;117.
- 152 Di Lorenzo R, Frattini N, Dragone D, *et al.* Psychiatric Emergencies During the Covid-19 Pandemic: A 6-Month Observational Study. *Neuropsychiatr Dis Treat* 2021;17:1763–78.
- 153 Schweiberger K, Patel SY, Mehrotra A, *et al.* Trends in Pediatric Primary Care Visits During the Coronavirus Disease of 2019 Pandemic. *Acad Pediatr* 2021;21:1426–33.
- 154 Peterson J, Welch Losos M, Tugwell P, *et al.* The newcastle-ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Hospital Research Institute; 2014.
- 155 Stroever S, Brett C, Michael K, *et al.* Emergency department utilization for mental health conditions before and after the COVID-19 outbreak. *Am J Emerg Med* 2021;47:164–8.
- 156 Fasshauer JM, Bollmann A, Hohenstein S, *et al.* Emergency hospital admissions for psychiatric disorders in a German-wide hospital network during the COVID-19 outbreak. *Soc Psychiatry Psychiatr Epidemiol* 2021;56:1469–75.
- 157 Enrico C, Caldiroli A, Di Brita C, *et al.* Profile of patients attending psychiatric emergency care during the coronavirus 2019 (COVID 19) pandemic: a comparative cross-sectional study between lockdown and post-lockdown periods in Lombardy, Italy. *Int J Psychiatry Clin Pract* 2022;26:132–8.
- 158 Clerici M, Durbano F, Spinogatti F, *et al.* Psychiatric hospitalization rates in Italy before and during COVID-19: did they change? An analysis of register data. *Ir J Psychol Med* 2020;37:283–90.
- 159 Simpson SA, Loh RM, Cabrera M, *et al.* The Impact of the COVID-19 Pandemic on Psychiatric Emergency Service Volume and Hospital Admissions. *J Acad Consult Liaison Psychiatry* 2021;62:588–94.
- 160 Pham-Scottet A, Silva J, Barruel D, *et al.* Patient flow in the largest French psychiatric emergency centre in the context of the COVID-19 pandemic. *Psychiatry Res* 2020;291:113205.

- 161 Hoyer C, Ebert A, Szabo K, *et al.* Decreased utilization of mental health emergency service during the COVID-19 pandemic. *Eur Arch Psychiatry Clin Neurosci* 2021;271:377–9.
- 162 Zielasek J, Vrinssen J, Gouzoulis-Mayfrank E. Utilization of Inpatient Mental Health Care in the Rhineland During the COVID-19 Pandemic. *Front Public Health* 2021;9:593307.
- 163 Finkelstein Y, Maguire B, Zemek R, *et al.* Effect of the COVID-19 Pandemic on Patient Volumes, Acuity, and Outcomes in Pediatric Emergency Departments. *Pediatr Emer Care* 2021;37:427–34.
- 164 Amerio A, Aguglia A, Odone A, *et al.* Covid-19 pandemic impact on mental health of vulnerable populations. *Acta Biomed Jul* 2020;91:95–6.
- 165 Mafham MM, Spata E, Goldacre R, *et al.* COVID-19 pandemic and admission rates for and management of acute coronary syndromes in England. *Lancet* 2020;396:381–9.
- 166 Nappi C, Megna R, Acampa W, *et al.* Effects of the COVID-19 pandemic on myocardial perfusion imaging for ischemic heart disease. *Eur J Nucl Med Mol Imaging* 2021;48:421–7.
- 167 Bonello F, Zammit D, Grech A, *et al.* Effect of COVID-19 pandemic on mental health hospital admissions: comparative population-based study. *BJPsych Open* 2021;7:e141.
- 168 Prina E, Tedeschi F, Salazzari D, *et al.* Effect of COVID-19 pandemic on utilisation of community-based mental health care in North-East of Italy: A psychiatric case register study. *Epidemiol Psychiatr Sci* 2023;32:e17.
- 169 White J, Cavenagh D, Byles J, *et al.* The experience of delayed health care access during the COVID 19 pandemic in Australian women: A mixed methods exploration. *Health Soc Care Community* 2022;30:e1384–95.
- 170 Shirzad F, Shati M, Mortazavi SS, *et al.* Outpatient psychiatric service utilization during the Covid-19 pandemic. *Middle East Curr Psychiatry* 2022;29.
- 171 Menges D, Ballouz T, Anagnostopoulos A, *et al.* Burden of post-COVID-19 syndrome and implications for healthcare service planning: A population-based cohort study. *PLoS One* 2021;16:e0254523.
- 172 Håkansson A, Grudet C. Decreasing Psychiatric Emergency Visits, but Stable Addiction Emergency Visits, During COVID-19-A Time Series Analysis 10 Months Into the Pandemic. *Front Psychiatry* 2021;12:664204.
- 173 Savić A, Vukojević J, Mitreković I, *et al.* Changes in psychiatric services dynamics during the COVID-19 pandemic: Recognizing the need for resources shift. *Psychiatry Res* 2022;317:114778.