

Original Article

*These authors contributed equally to this work.

Cite this article: Spinazzola E *et al* (2023). The association between reasons for first using cannabis, later pattern of use, and risk of first-episode psychosis: the EU-GEI case-control study. *Psychological Medicine* 1–10. <https://doi.org/10.1017/S0033291723001071>

Received: 21 September 2022

Revised: 23 January 2023

Accepted: 3 April 2023




Keywords:

Cannabis use; path analysis; psychotic disorders

Corresponding author:

Edoardo Spinazzola,
E-mail: Edoardo.spinazzola@kcl.ac.uk

The association between reasons for first using cannabis, later pattern of use, and risk of first-episode psychosis: the EU-GEI case-control study

Edoardo Spinazzola^{1,*} , Diego Quattrone^{2,3,4,*} , Victoria Rodriguez¹, Giulia Trotta² , Luis Alameda^{1,5,6} , Giada Tripoli^{1,7} , Charlotte Gayer-Anderson⁸ , Tom P Freeman^{9,10}, Emma C Johnson¹¹, Hannah E Jongsma¹², Simona Stilo^{1,13}, Caterina La Cascia⁷, Laura Ferraro⁷, Daniele La Barbera⁷, Antonio Lasalvia¹⁴, Sarah Tosato¹⁴, Iliaria Tarricone¹⁵, Giuseppe D'Andrea¹⁵, Michela Galatolo¹⁵, Andrea Tortelli^{16,17}, Iliaria Tagliabue^{18,19}, Marco Turco¹⁸, Maurizio Pompili²⁰, Jean-Paul Selten^{21,22}, Lieuwe de Haan²³, Paulo Rossi Menezes²⁴, Cristina M Del Ben²⁴, Jose Luis Santos²⁵, Manuel Arrojo²⁶, Julio Bobes²⁷, Julio Sanjuán²⁸, Miguel Bernardo²⁹, Celso Arango³⁰, James B Kirkbride³¹, Peter B Jones^{32,33}, Michael O'Donovan³⁴, Bart P Rutten²², Jim Van Os^{1,21,35}, Craig Morgan⁸, Pak C Sham³⁶, Isabelle Austin-Zimmerman², Zhikun Li², Evangelos Vassos², EU-GEI WP2 Group¹, Robin M Murray¹ and Marta Di Forti^{2,4,37} 

Abstract

Background. While cannabis use is a well-established risk factor for psychosis, little is known about any association between reasons for first using cannabis (RFUC) and later patterns of use and risk of psychosis.

Methods. We used data from 11 sites of the multicentre European Gene-Environment Interaction (EU-GEI) case-control study. 558 first-episode psychosis patients (FEPp) and 567 population controls who had used cannabis and reported their RFUC. We ran logistic regressions to examine whether RFUC were associated with first-episode psychosis (FEP) case-control status. Path analysis then examined the relationship between RFUC, subsequent patterns of cannabis use, and case-control status.

Results. Controls (86.1%) and FEPp (75.63%) were most likely to report 'because of friends' as their most common RFUC. However, 20.1% of FEPp compared to 5.8% of controls reported: 'to feel better' as their RFUC ($\chi^2 = 50.97$; $p < 0.001$). RFUC 'to feel better' was associated with being a FEPp (OR 1.74; 95% CI 1.03–2.95) while RFUC 'with friends' was associated with being a control (OR 0.56; 95% CI 0.37–0.83). The path model indicated an association between RFUC 'to feel better' with heavy cannabis use and with FEPp-control status.

Conclusions. Both FEPp and controls usually started using cannabis with their friends, but more patients than controls had begun to use 'to feel better'. People who reported their reason for first using cannabis to 'feel better' were more likely to progress to heavy use and develop a psychotic disorder than those reporting 'because of friends'.

Introduction

A meta-analysis including cross-sectional and longitudinal studies has shown that heavy cannabis use is associated with a 4-fold increase in the risk of psychosis (Marconi, Di Forti, Lewis, Murray, & Vassos, 2016). Moreover, a higher incidence of psychotic disorders was reported in regions with a higher prevalence of daily use and greater availability of high-potency cannabis (Di Forti *et al.*, 2019). The average proportion of delta-9-tetrahydrocannabinol (THC) in the cannabis available in international markets increased from 1970 to 2017 (Freeman *et al.*, 2021). Two studies conducted in Denmark found that an increase in the use and potency of cannabis in the population during the past two decades was accompanied by a noticeable increase in the incidence of cannabis-induced psychosis (Hjorthoj, Larsen, Starzer, & Nordentoft, 2021a), and a 3- to 4-fold increase in the proportion of cases of schizophrenia associated with cannabis use disorder (CUD) (Hjorthoj, Posselt, & Nordentoft, 2021b). Moreover, a recent within-person

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

analysis – where each participant serves as their control – found that cannabis is likely to have a causal impact on psychosis, but not the other way around (van Os et al., 2021).

Some argue that the association between cannabis use and psychotic disorders can be explained as an attempt to ‘self-medicate’ by those who experience sub-clinical psychosis, trying to relieve existing subthreshold psychotic symptoms (Khantzian, 1985). However, most studies report that patients with psychosis use cannabis for the same reasons as the general population, such as for the pleasurable effects and social reasons rather than to self-medicate (Dekker, Linszen, & De Haan, 2009; Dekker et al., 2010; Green, Kavanagh, & Young, 2004; Kolliakou, Joseph, Ismail, Atakan, & Murray, 2011; Pérez, Santacana, Baquero, & Pérez-Solà, 2014).

While some studies have examined why people use cannabis and whether the subjective effects of cannabis differ between patients with their first episode of psychosis and controls (Bianconi et al., 2016; Dekker, Koeter, Van Den Brink, & Investigators, 2012; Dekker et al., 2009; Peters et al., 2009), an accurate search of the literature (see online supplementary) indicated little evidence on the reasons for the first use of cannabis (RFUC). Moreover, it is not clear if RFUC explains later patterns of use, or if RFUC differs between those who later develop psychosis and healthy controls.

This is the first study that uses data from a large multisite first-episode psychosis (FEP) case-control design and aims to examine (1) the socio-demographic factors associated with RFUC and (2) which RFUC, among cannabis users, is associated with harmful patterns of cannabis use and with an increased risk for psychotic disorders.

Methods

Sample

This investigation is based on the European network of national schizophrenia networks studying Gene-Environment Interactions (EU-GEI study, <http://www.eu-gei.eu>), a multicentre incidence and case-sibling-control study of genetic and environmental determinants of psychotic disorder (Gayer-Anderson et al., 2020; Jongsma et al., 2018) consisting of first-episode psychosis patients (FEPp) and population-based controls recruited between 2010 and 2015, who provided information on cannabis use (Gayer-Anderson et al., 2020). Both FEPp and controls were collected in 17 catchment areas in six countries: Southeast London, Cambridgeshire and Peterborough (England); central Amsterdam, Gouda and Voorhout (the Netherlands); part of the Veneto region, Bologna municipality, city of Palermo (Italy); 20th Arrondissement of Paris, Val-de-Marne, Puy-de-Dôme (France); Madrid (Vallecas), Barcelona, Valencia, Oviedo, Santiago, Cuenca (Spain); and Ribeirão Preto (Brazil).

Participants

All patients presenting with FEP who were referred to the mental healthcare services within the 17 catchment areas were identified by trained researchers. Patients were included in the current study if they were aged 18–64 years and resident within the study areas at the time of their first presentation and if they presented with a clinical diagnosis for an untreated FEP (*International Statistical Classification of Diseases, Tenth Revision (ICD-10)* codes F20–F33). Exclusion criteria were: (a) previous contact with psychiatric services for psychosis; (b) evidence of psychotic symptoms precipitated by an organic cause; and (c) transient psychotic symptoms

resulting from acute intoxication, as defined by the ICD-10 (codes F1X.5). Controls were recruited using quota sampling based on local census data to ensure samples representativeness in each catchment area’s population at risk in terms of age, gender, and ethnicity. Inclusion criteria for controls were: (a) aged between 18 and 64 years; (b) resident within the catchment areas at the time of consent into the study; (c) sufficient command of the primary language at each site to complete the assessments; and (d) no current or past psychotic disorder. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All participants who agreed to take part in the study provided informed, written consent; and ethical approval was provided by research ethics committees in each site: South London and Maudsley and Institute of Psychiatry Research Ethics Committee; National Research Ethics Service Committee East of England–East Cambridge; Medisch-Ethische Toetsingscommissie van het Academisch Centrum te Amsterdam; Comité Ético de Investigación Clínica Hospital Gregorio Marañón; Comité Ético de Investigación Clínica del Hospital Clinic de Barcelona; Comité Ético de Investigación Clínica del Hospital Clinic Universitari de Valencia; Comité Ética de la Investigación Clínica del Principado de Asturias; Comité Ético de Investigación Clínica de Galicia; Comité Ético de Investigación Clínica del Hospital Virgen de la Luz de Cuenca; Comité de Protección des Personnes–CPP Île de France IX; Comitato Etico Policlinico S Orsola Malpighi; Comitato Etico Azienda Ospedaleria Universitaria di Verona; Comitato Etico Palermo 1, Azienda Ospedaliere Policlinico ‘Paolo Giaccone’; and Research Ethics Committee of the clinical Hospital of Ribeirão Preto Medical School, University of São Paulo, Brazil. Further information on the general study methods is available in the EU-GEI core paper on the description of the objectives and main aspects of the study (Gayer-Anderson et al., 2020).

Measures and assessments

Data on age, gender, and self-reported ethnicity were collected using a modified version of the Medical Research Council Sociodemographic Schedule (Mallett, Leff, Bhugra, Pang, & Zhao, 2002). Diagnoses of psychotic disorders were confirmed using the OPERational CRITERia (OPCRIT) system (McGuffin, Farmer, & Harvey, 1991; Williams, Farmer, Ackenheil, Kaufmann, & McGuffin, 1996), which was completed by centrally trained investigators whose inter-rater reliability was assessed throughout the study ($\kappa = 0.7$). Family history of mental illness and parental history of psychosis were collected using the Family Interview for Genetic Studies questionnaire (Maxwell, 1992). Finally, the Intelligence Quotient (IQ) score was derived using the short form of the WAIS-III (IQ), including selected items of the following subtests: digit symbol coding (a measure of processing speed), arithmetic (working memory), block design (visuospatial processing), and information (verbal knowledge) (Tripoli et al., 2021).

Detailed information on cannabis use was collected using the latest version of the Cannabis Experiences Questionnaire (Barkus, Stirling, Hopkins, & Lewis, 2006), for the EU-GEI study (CE_{EU-GEI}). Consistently with Di Forti et al. (2019), we used the combined measure of frequency and type of cannabis used derived from the following variables: (1) Frequency of use: never or occasional = 0, more than once a week = 1, daily = 2; (2) low potency cannabis with less than 10% of THC = 0, high potency THC 10%




Figure 1. Overlapping reported RFUC in cases and controls.

or greater = 1 (see online supplementary materials for a detailed description of this variable). This combined measure was used to define the *harmful pattern of use* and coded as follows: occasional use with any potency of use = 1, more than once a week and low potency = 2, more than once a week and high potency = 3, daily or almost and low potency = 4, daily or almost and high potency = 5 (see online supplementary). Age at first cannabis use was considered as a continuous numerical variable.

The information on RFUC was derived from the responses to the question ‘why did you first try cannabis?’ in the form of the following multiple-choice: (a) my friends were using it; (b) my family members were using it; (c) to feel better (to get relief from either physical or psychological discomfort) and (d) other. Subjects were able to provide up to 4 RFUC.

Statistical analysis

We compared the number of different reported RFUC in cases and controls, also taking into account the overlapping answers.

T-test (Mann–Whitney where appropriate) and χ^2 in STATA16 were used to test for association between socio-demographics and IQ with RFUC. Only the variables associated with RFUC were included as covariates in the Path analyses.

We conducted unadjusted and adjusted logistic regressions using STATA-16 to examine whether RFUC was associated with FEP case–control status, adjusting for IQ score, gender, age at first cannabis use, ethnic minority status, and the harmful pattern of cannabis use. We conducted the analysis in stages to test four models. Model A examined the unadjusted associations between different RFUC and FEP case–control status. Model B examined the associations after adjusting for the harmful patterns of use. Model C additionally adjusted for age at first cannabis use. Lastly, model D additionally adjusted for ethnic minority status, IQ score, and gender. We compared the statistical fit of the four

models using Log-Likelihood (LL), Akaike information criteria (AIC), and Bayesian information criteria (BIC).

Path analysis (Stage, Carter, & Nora, 2004) was run in STATA-16 to test the hypothesised direction in the relationship between multiple RFUC as predictors, with (1) pattern of cannabis use and (2) FEP case–control status.

In the regression analyses and in the path model, we included the participants who reported to have used cannabis either because of ‘friends’, ‘family’, or ‘to feel better’.

Results

Socio-demographic characteristics

Participants were recruited and consented to the study between May 1, 2010, and April 1, 2015. The 20th arrondissement of Paris was the only one that did not contribute to the recruitment of population controls; hence, it was excluded from the analyses. As in a previous EU-GEI paper (Di Forti et al., 2019), we also excluded Verona, Santiago, Oviedo, Valencia, and Cuenca because they had at least 10% of data missing on the measures of cannabis use. This left 901 FEPp and 1235 controls (see recruitment flow chart in the online supplementary materials).

585 FEPp (64.93%) and 574 (46.4%) of the controls reported lifetime cannabis use. Among these, 27 FEPp (4.62%) and 7 controls (1.22%) were excluded because of missing data on RFUC. Therefore, the sample resulted in a total size of 558 FEPp and 567 controls.

We found that, among users, 488 (86.1%) controls and 422 (75.63%) FEPp reported RFUC because of ‘friends’, while 38 (6.70%) controls and 69 (12.37%) FEPp reported RFUC because of their ‘family’ members; 33 (5.82%) controls and 112 (20.07%) of FEPp started ‘to feel better’, and 159 (28.04%) controls and 218 (39.07%) of FEPp because of ‘other’ reasons. See Fig. 1 for detailed information on the overlapping answers for this variable.

Cannabis users who reported RFUC because of ‘friends’ did not show any association with the socio-demographic variables assessed. However, those who reported RFUC ‘to feel better’ were more often from ethnic minorities ($\chi^2(1) = 13.67$; $p < 0.001$) and had lower IQ scores ($U = 6.66$; $p < 0.001$). Similarly, those who reported RFUC because of their ‘family’ members, were more often from ethnic minorities ($\chi^2(1) = 7.97$; $p = 0.005$) and had lower IQ scores ($U = 3.11$; $p = 0.002$). Lastly, those who reported RFUC ‘other’ reasons were more likely to be male ($\chi^2(1) = 8.65$; $p = 0.003$) and were more often from ethnic minorities ($\chi^2(1) = 4.43$; $p = 0.035$; Table 1).

FEPp were more likely to belong to ethnic minority groups ($\chi^2(1) = 62.37$; $p < 0.001$) and had lower IQ ($U = 14.97$; $p < 0.001$) compared to controls (see online supplementary materials).

Logistic regressions

In the regression and in the path models, we included the participants who reported that they had used cannabis either because of ‘friends’, ‘family’, or ‘to feel better’, thus excluding 59 controls (10.41%) and 83 cases (14.87%) who reported ‘other’ as their RFUC. This resulted in a final sample of 475 FEPp and 508 controls with a complete dataset on reasons to start using cannabis.

The Model A unadjusted logistic regression indicated that RFUC because of ‘friends’ was negatively associated with being a FEP (OR 0.58; 95% CI 0.43–0.79), while those reporting RFUC because of ‘family’ members (OR 1.59; 95% CI 1.03–2.44) or ‘to feel better’ (OR 3.86; 95% CI 2.56–5.82) were more likely to become FEPp. In Model B, after controlling for the harmful patterns of use, RFUC because of ‘family’ members was no longer associated with FEP case–control status, while RFUC because of ‘friends’ was still associated with being a control (OR 0.64; 95% CI 0.45–0.91) and RFUC ‘to feel better’ was still associated with being a FEPp (OR 2.60; 95% CI 1.67–4.04). In Model C, after additionally adjusting for age at first cannabis use, the results remained consistent with Model B for both RFUC because of ‘friends’ (OR 0.62; 95% CI 0.44–0.88) and ‘to feel better’ (OR 2.63; 95% CI 1.69–4.10). In Model D, additionally adjusting for ethnic minority status, IQ score, and gender, RFUC because of ‘friends’ (OR 0.56; 95% CI 0.37–0.83) and ‘to feel better’ (OR 1.74; 95% CI 1.03–2.95) were still associated with FEP case–control status (See Table 2).

Path analysis

The path model had a good fit to the data: root mean square error of approximation (RMSEA) = 0.040, comparative fit index (CFI) = 0.99 (Fig. 2). RFUC because of ‘friends’ was associated with being a control ($\beta = -0.08$; $p = 0.002$), while ‘to feel better’ was associated with being a FEPp ($\beta = 0.07$; $p = 0.014$) (see Table 3 for direct associations). Lower IQ scores were indirectly associated with RFUC ‘to feel better’ ($\beta = -0.19$; $p \leq 0.001$), with a younger age at first cannabis use ($\beta = 0.11$; $p = 0.001$), harmful pattern of cannabis use ($\beta = -0.16$; $p \leq 0.001$) and were directly associated with being a FEPp ($\beta = -0.36$; $p \leq 0.001$). On the contrary, higher IQ scores were associated with RFUC because of ‘friends’ ($\beta = 0.07$; $p = 0.043$), which was negatively associated with age at first cannabis use ($\beta = -0.10$; $p = 0.002$).

Both RFUC because of ‘family’ ($\beta = 0.09$; $p = 0.003$) and ‘to feel better’ ($\beta = 0.15$; $p \leq 0.001$) were associated with a more harmful pattern of cannabis use, the latter also with being a FEPp. See Table 3 for direct and indirect associations.

Table 1. Descriptive sociodemographic analyses based on reported RFUC (in cannabis users)

		‘Friends’ yes/no N (% or SD)	Statistics and p value	‘Family’ yes/no N (% or SD)	Statistics and p value	‘To feel better’ yes/no N (% or SD)	Statistics and p value	‘Other’ yes/no N (% or SD)	Statistics and p value
Gender	Male	560 (61.54)	$\chi^2 = 0.84$ $p = 0.36$	71 (66.36)	$\chi^2 = 0.86$ $p = 0.35$	97 (66.90)	$\chi^2 = 1.58$ $p = 0.21$	257 (68.17)	$\chi^2 = 8.65$ $p = 0.003$
Ethnic minority status	Yes	254 (27.94)	$\chi^2 = 0.49$ $p = 0.48$	43 (40.19)	$\chi^2 = 7.97$ $p = 0.005$	60 (41.38)	$\chi^2 = 13.67$ $p < 0.001$	122 (32.45)	$\chi^2 = 4.43$ $p = 0.035$
Ethnicity	White	655 (72.06)	$\chi^2 = 2.96$ $p = 0.57$	64 (59.81)	$\chi^2 = 15.88$ $p = 0.003$	85 (58.62)	$\chi^2 = 17.38$ $p = 0.002$	254 (67.55)	$\chi^2 = 10.41$ $p = 0.034$
	Black	115 (12.65)		21 (19.63)		26 (17.93)		48 (12.77)	
	Mixed	71 (7.81)		16 (14.95)		21 (14.48)		42 (11.17)	
	Asian	22 (2.42)		1 (0.93)		4 (2.76)		13 (3.46)	
	Other	46 (5.06)		5 (4.67)		9 (6.21)		19 (5.05)	
IQ		97.30 (20.25)	$U = -1.95$ $p = 0.051$	91.08 (19.89)	$U = 3.11$ $p = 0.002$	86.10 (17.25)	$U = 6.66$ $p < 0.001$	96.35 (19.15)	$U = 0.59$ $p = 0.56$
Parental mental illness	Yes	310 (34.33)	$\chi^2 = 0.02$ $p = 0.88$	42 (39.25)	$\chi^2 = 1.28$ $p = 0.26$	51 (35.66)	$\chi^2 = 0.12$ $p = 0.73$	132 (35.58)	$\chi^2 = 0.36$ $p = 0.55$
Parental Psychosis	Yes	64 (7.09)	$\chi^2 = 0.68$ $p = 0.41$	11 (10.28)	$\chi^2 = 1.54$ $p = 0.21$	14 (9.79)	$\chi^2 = 1.38$ $p = 0.24$	31 (8.36)	$\chi^2 = 0.75$ $p = 0.39$

χ^2 , Chi-squared test; U, Mann–Whitney U test; *p-value ≤ 0.05 .

Table 2. Unadjusted and adjusted associations between RFUC and case-control status

	Case-control outcome			
	Model A R (95% CI)	Model B Adjustment for harmful pattern of use OR (95% CI)	Model C Plus adjustment for age at first cannabis use	Model D Plus adjustment for ethnic minority status and IQ score
Friends	0.58 (0.43–0.79)	0.64 (0.45–0.91)	0.62 (0.44–0.88)	0.56 (0.37–0.83)
Family	1.59 (1.03–2.44)	1.12 (0.69–1.82)	1.12 (0.69–1.81)	0.99 (0.57–1.73)
Better	3.86 (2.56–5.82)	2.60 (1.67–4.04)	2.63 (1.69–4.10)	1.74 (1.03–2.95)

Model A: unadjusted associations; Model B: adjusted for harmful pattern of use; Model C: adjusted for harmful pattern of use and age at first cannabis use; Model D: adjusted for harmful pattern of use, age at first cannabis use, ethnic minority status, and IQ score.




Figure 2. Direct and indirect pathways between IQ, ethnicity, RFUC and case-control status. Significant pathways signified by solid arrows and *; nonsignificant pathways represented by dotted lines. Model fit: $\chi^2 = 12.50$, RMSEA = 0.040, comparative fit index CFI = 0.99..

Discussion

Principal findings

To the best of our knowledge (see online supplementary materials for details of literature search), this is the first study to examine what reasons underlie first using cannabis and if these reasons are associated with later pattern of cannabis use and the risk to develop a psychotic disorder.

First, our findings indicate that having friends who use cannabis is the most common reason to start using cannabis among both those with FEPP and the controls; although, a higher proportion of FEPP than controls reported ‘to feel better’ as their RFUC (Fig. 1). Second, we provide the first evidence that, (a) compared to those who RFUC with ‘friends’, those who reported starting ‘to feel better’ as their RFUC are much more likely to progress to daily use of high potency cannabis; and (b) those who started to use cannabis because of their ‘friends’ are more likely to have started their use earlier in their life (Table 3).

Limitations and strengths

Our findings need to be appraised in the context of some limitations. Firstly, the data on cannabis use were collected retrospectively based on self-report. Therefore, they may be open to recall bias. However, studies comparing laboratory data and self-reported information have shown that cannabis users reliably report their frequency of use as well as the type of cannabis they use

(Buchan, Dennis, Tims, & Diamond, 2002; Freeman et al., 2014). While it is possible that FEPP are more likely, compared to controls, to retrospectively report ‘to feel better’ as a RFUC to justify their daily use leading to recall bias, the category ‘to feel better’ includes relief from both physical and psychological discomfort, therefore it is not intended to capture self-medication from prodromal symptoms. Moreover, a previous study has shown that cannabis use is associated with increased risk of psychosis even after adjustment for baseline prodromal symptoms (Mustonen et al., 2018). A recent study pointed out that some people might use cannabis to ameliorate anxiety and depressive symptoms (Radhakrishnan et al., 2022). However, the evidence suggests that cannabis use may increase the risk of developing depression and suicidality (Gobbi et al., 2019). A recent large online survey including responses from 27 169 participants from Canada and the USA, showed that 53% of people using cannabis to relieve physical discomfort reported using it to alleviate pain, followed by difficulties with sleep and headache/migraine as reasons to use (Leung et al., 2022). These are physical complaints persisting in nature and it is plausible that the attempt to alleviate them might lead to regular rather than occasional use.

Thirdly, although pathway analysis tests causal inferences and linkages among variables in the context of an a-priori conceptual model, causation cannot be implied. Despite the validity of path analysis in cross-sectional studies has been confirmed (Etain et al., 2017; Kwok, Cheung, Jak, Ryu, & Wu, 2018; Martin, 2011), prospective longitudinal studies will be required to confirm

Table 3. Panel A: Standardised Probit Coefficients (β) for the indirect pathways to psychosis with RFUC and pattern of cannabis use as mediators; Panel B Standardised Probit Coefficients (β) for the direct pathways to psychosis

PANEL A. Indirect pathways to psychosis with RFUC and pattern of cannabis use as mediators															
Predictor variables	Via Friends			Via Family			Via to feel better			Via Age at first cannabis use			Via Frequency-potency of cannabis use		
	β	s.e.	p	β	s.e.	p	β	s.e.	p	β	s.e.	p	β	s.e.	p
IQ scores	0.07	0.03	0.043	-0.06	0.04	0.066	-0.19	0.03	<0.001	0.11	0.03	0.001	-0.16	0.03	<0.001
Ethnic minority Status	N/A	N/A	N/A	-0.05	0.03	0.12	0.03	0.03	0.336	N/A	N/A	N/A	0.07	0.03	0.025
Friends	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	-0.10	0.03	0.002	-0.09	0.03	0.002
Family	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	-0.02	0.03	0.49	0.09	0.03	0.003
To feel better	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	-0.04	0.03	0.209	0.15	0.03	<0.001
PANEL B. Direct pathways to psychosis															
Predictor variables	Case-control status														
	β			s.e.			p								
IQ scores	-0.36			0.03			<0.001								
Ethnic minority status	0.07			0.03			0.017								
Friends	-0.08			0.03			0.002								
Family	-0.003			0.03			0.913								
To feel better	0.07			0.03			0.014								
Age at first cannabis use	-0.03			0.03			0.324								
Frequency-potency of cannabis use	0.25			0.29			<0.001								

Note: β , Probit coefficient; SE, Standard error; p, probability; N/A, Not applicable.

the detected associations, which nevertheless, in our study clearly places first use of cannabis on average at least a decade before the psychosis onset. Furthermore, this is a FEPp-control study which is less likely to lead to recall bias because the participants are recruited near the onset of their psychosis illness.

Finally, the data presented here were collected from a control sample representative of each site's local population at risk (Gayer-Anderson et al., 2020) and from a subset of FEPp representative of the larger incidence sample recruited from each site Mental Health services over the study period (Jongsma et al., 2018). Thus, our findings have the important strength of describing the RFUC reported by individuals with and without psychosis, from different ethnicities across different geographical areas and are therefore generalisable. Moreover, the use of matched controls might have masked the association between sociodemographic variables with RFUC and pattern of cannabis use.

Comparison with previous research

Previous research indicated that earlier age at first use is more likely to progress to longstanding use, resulting in overall greater exposure to cannabis (Radhakrishnan, Wilkinson, & D'Souza, 2014). Furthermore, data have suggested that individuals who start using cannabis early in adolescence may be at most risk to develop psychotic disorder (Arseneault et al., 2002; Dragt et al., 2010; Korver et al., 2010); one study suggested they might 'self-medicate' with cannabis to alleviate initial symptoms (Ferdinand et al., 2005). In contrast, our findings do not show an association between RFUC to 'feel better' and starting to use cannabis early in adolescence, which might be explained by our category 'to feel better' referring to relief not only of psychological but also physical discomfort. A recent study, in fact, shows how cannabis use for medical reasons is more common among young adults and older age groups rather than adolescence (Leung et al., 2022). Instead, we found that those who reported a younger age at first cannabis use were more likely to have started using cannabis 'with friends' compared to the other groups. This is in line with previous evidence looking at reasons to use cannabis which reports social context as the main reason for use (Koliakou et al., 2011). The high number of cannabis users who start 'with friends' is in accord with previous findings indicating that cannabis users with psychosis have better premorbid social functioning compared to patients with psychosis not using cannabis (Ferraro et al., 2021; Ferraro et al., 2020; Ferraro et al., 2013). A possible explanation for this is that cannabis-using patients with psychosis are more socially skilled, and therefore able to obtain the substance than those who are neurodevelopmentally impaired (Murray et al., 2017). We found no evidence that those who report RFUC 'to feel better' are more likely to start using cannabis close to their age of psychosis onset.

RFUC because of 'family members' is associated with a more harmful pattern of use. This suggests that targeting the family environment could play an important role in delivering interventions for substance misuse and also to disseminate education messages about the risk associated with cannabis use and its harmful effects on mental health.

The existing evidence clearly indicates that daily use especially of high potency cannabis is robustly associated with an increased probability to develop a psychotic disorder (Di Forti et al., 2009, 2015, 2019). While we cannot separate those seeking relief from physical or psychological discomfort, our data suggest that starting to use cannabis 'to feel better' is more likely to progress to

daily use of high potency cannabis and to later suffer a first episode psychosis.

In conclusion, while only a minority of cannabis users develops psychosis, understanding what drives people to first use cannabis can provide valuable data on (a) how to identify those more likely to develop a harmful pattern of use and design interventions for harm reduction or use cessation tailored to the individual's needs, (b) provide support and close monitoring to those using cannabis for medical reasons to minimise the risks associated with regular use, and (c) design better public health campaigns that are able to reach the individual in the social and family context where first use is most likely to begin.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291723001071>.

Data. The data that support the findings of the study is available on request from the corresponding author, E.S.

Author contributions. E. S., D. Q., V. R., M. D. F., and R. M. contributed to the conceptualization and design of the study. E. S., D. Q., M. D. F., and R. M. took part in the interpretation of the data and wrote the first draft. All other authors contributed to the revision of the manuscript. They also read and approved the final manuscript.

Financial support. This study was funded by the Medical Research Council, the European Community's Seventh Framework Program grant (agreement HEALTH-F2-2009-241909 (Project EU-GEI), Sao Paulo Research Foundation (grant 2012/0417-0), the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) at South London and Maudsley NHS Foundation Trust and King's College London, the NIHR BRC at University College London, and the Wellcome Trust (grant 101272/Z/12/Z). ES was supported by Lord Leverhulme's Charitable Trust and the Velvet Foundation and by the Medical Research Council (MRC) – UKRI [MR/T007818/1]. MDF and GT are supported by the Medical Research Council (MRC) – UKRI [MR/T007818/1]. DQ is supported by MRC/UKRI CARP (MRC CARP grant MR/W030608/1). Dr Kirkbride is supported by the National Institute for Health Research (NIHR) University College London (UCL) Biomedical Research Centre (BRC). Dr Emma Johnson receives funding from NIDA: K01DA051759.

Conflict of interest. J Bobes has received research grants and served as consultant, advisor or speaker for AB-Biotics, Acadia Pharmaceuticals, Ambrossetti-Angelini, Casen Recordati, D&A Pharma, Exeltis, Gilead, Indivior, Janssen-Cilag, Lundbeck, Mundipharma, Otsuka, Pfizer, Roche, Sage Therapeutics, Servier, Schwabe Farma Ibérica, Shire, Takeda, research funding from the Spanish Ministry of Economy and Competitiveness – Centro de Investigación Biomedica en Red area de Salud Mental (CIBERSAM) and Instituto de Salud Carlos III-, Spanish Ministry of Health, Social Services and Equality - Plan Nacional sobre Drogas outside of the submitted work. M Bernardo has been a consultant for, received grant/research support and honoraria from, and been on the speakers/advisory board of AB-Biotics, Adamed, Angelini, Casen Recordati, Janssen-Cilag, Menarini, Rovi and Takeda. C Arango reports personal fees from Acadia, Ambrossetti, Gedeon Richter, Janssen Cilag, Lundbeck, Merck, Otsuka, Roche, Servier, Shire, Schering Plough, Sumitomo Dainippon Pharma, Sunovion, and Takeda; and grants from CIBERSAM, Familia Alonso, Fundacion Alicia Koplowitz, the European Commission, the Spanish Ministry of Science and Universities, and the Comunidad de Madrid, during the conduct of the study. PB Jones reports personal fees from being a member of the scientific advisory boards for Janssen and Recordati, outside of the submitted work. Prof Michael O' Donovan is supported by a collaborative research grant from Takeda outside of the submitted work. R M Murray reports personal fees from Janssen, Lundbeck, Sunovion, and Otsuka, outside of the submitted work. M Di Forti reports personal fees from Janssen, outside the submitted work. All other authors declare no competing interests.

EU-GEI Collaborators. Kathryn Hubbard¹, Stephanie Beards¹, Simona A. Stilo², Mara Parellada³, David Fraguas³, Marta Rapado Castro³, Álvaro

Andreu-Bernabeu³, Gonzalo López³, Mario Matteis³, Emiliano González³, Manuel Durán-Cutilla³, Covadonga M. Díaz-Caneja³, Pedro Cuadrado⁴, José Juan Rodríguez Solano⁵, Angel Carracedo⁶, Javier Costas⁶, Emilio Sánchez⁷, Silvia Amoretti⁸, Esther Lorente-Rovira⁹, Paz Garcia-Portilla¹⁰, Estela Jiménez-López¹¹, Eva Velthorst¹², Nathalie Franke¹², Daniella van Dam¹², Fabian Termorshuizen^{13,14}, Nathalie Franke¹³, Elsje van der Ven^{13,14}, Elles Messchaert¹⁴, Marion Leboyer^{15,16,17,18}, Franck Schürhoff^{15,16,17,18}, Stéphane Jamain^{16,17,18}, Grégoire Baudin^{15,16}, Aziz Ferchiou^{15,16}, Baptiste Pignon^{15,16,18}, Jean-Romain Richard^{16,18}, Thomas Charpeaud^{18,19,20}, Anne-Marie Tronche^{18,19,20}, Pierre-Michel Llorca²⁰, Flora Frijda^{21,22}, Giovanna Marrazzo²², Lucia Sideli²², Crocettarachele Sartorio^{22,23}, Fabio Seminerio²², Camila Marcelino Loureiro^{24,25}, Rosana Shuhama^{24,25}, Mirella Ruggeri²⁶, Chiara Bonetto²⁶, Dorian Cristofalo²⁶, Domenico Berardi²⁷, Marco Serri²⁷, Alexander Richards²⁸, and Aoife Coghlan²⁹.

¹Department of Health Service and Population Research, Institute of Psychiatry, Kings College London, De Crespigny Park, Denmark Hill, London SE5 8AF, UK.

²Department of Psychosis Studies, Institute of Psychiatry, King's College London, De Crespigny Park, Denmark Hill, London SE5 8AF, UK.

³Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), C/Doctor Esquerdo 46, 28007 Madrid, Spain. ⁴Villa de Vallecas Mental Health Department, Villa de Vallecas Mental Health Centre, Hospital Universitario Infanta Leonor / Hospital Virgen de la Torre, C/San Claudio 154, 28038 Madrid, Spain.

⁵Puente de Vallecas Mental Health Department, Hospital Universitario Infanta Leonor / Hospital Virgen de la Torre, Centro de Salud Mental Puente de Vallecas, C/Peña Gorbea 4, 28018 Madrid, Spain.

⁶Fundación Pública Galega de Medicina Xenómica, Hospital Clínico Universitario, Choupana s/n, 15782 Santiago de Compostela, Spain.

⁷Department of Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), C/Doctor Esquerdo 46, 28007 Madrid, Spain.

⁸Department of Psychiatry, Hospital Clinic, IDIBAPS, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Universidad de Barcelona, C/Villarroel 170, escalera 9, planta 6, 08036 Barcelona, Spain.

⁹Department of Psychiatry, School of Medicine, Universidad de Valencia, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), C/Avda. Blasco Ibáñez 15, 46010 Valencia, Spain.

¹⁰Department of Medicine, Psychiatry Area, School of Medicine, Universidad de Oviedo, Centro de

Investigación Biomédica en Red de Salud Mental (CIBERSAM), C/Julían Clavería s/n, 33006 Oviedo, Spain. ¹¹Department of Psychiatry, Servicio de Psiquiatría Hospital "Virgen de la Luz", C/Hermandad de Donantes de Sangre, 16002 Cuenca, Spain.

¹²Department of Psychiatry, Early Psychosis Section, Academic Medical Centre, University of Amsterdam,

Meibergdreef 5, 1105 AZ Amsterdam, The Netherlands.

¹³Rivierduinen Centre for Mental Health, Leiden, Sandiforddreef 19, 2333 ZZ Leiden, The Netherlands. ¹⁴Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, P.O. Box 616, 6200 MD Maastricht, The Netherlands.

¹⁵AP-HP, Groupe Hospitalier "Mondor", Pôle de Psychiatrie, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France.

¹⁶INSERM, U955, Equipe 15, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France.

¹⁷Faculté de Médecine, Université Paris-Est, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France.

¹⁸Fondation Fondamental, 40 Rue de Mesly, 94000 Créteil, France.

¹⁹CMP B CHU, BP 69, 63003 Clermont Ferrand, Cedex 1, France.

²⁰Université Clermont Auvergne, EA 7280, Clermont-Ferrand 63000, France.

²¹Etablissement Public de Santé Maison Blanche, Paris, France.

²²Department of Experimental Biomedicine and Clinical Neuroscience, Section of Psychiatry, University of Palermo, Via G. La Loggia n.1, 90129 Palermo, Italy.

²³Unit of Psychiatry, "P. Giaccone" General Hospital, Via G. La Loggia n.1, 90129 Palermo, Italy. ²⁴Departamento de Neurociências e Ciências do Comportamento, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Av. Bandeirantes, 3900 -Monte Alegre- CEP 14049-900, Ribeirão Preto, SP, Brasil.

²⁵Núcleo de Pesquisa em Saúde Mental Populacional, Universidade de São Paulo, Avenida

Doutor Arnaldo 455, CEP 01246-903, Ribeirão Preto, SP, Brasil.

²⁶Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Piazzale L. A. Scuro 10, 37134 Verona, Italy.

²⁷Department of Medical and Surgical Science, Psychiatry Unit, Alma Mater Studiorum Università di Bologna, Viale Pepoli 5, 40126 Bologna, Italy.

²⁸Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, UK.

²⁹Family Genomics Research Group, Department of Biology, Maynooth University, Maynooth, Ireland.

¹Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College of London, London, UK; ²Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE5 8AF, UK; ³National Institute for Health Research, Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College, London, UK;

⁴South London and Maudsley NHS Mental Health Foundation Trust, London, UK; ⁵Service of General Psychiatry, Treatment and Early Intervention in Psychosis Program, Lausanne, University Hospital (CHUV), Lausanne, Switzerland; ⁶Centro Investigación Biomédica en Red de Salud Mental (CIBERSAM); Instituto de Biomedicina de Sevilla (IBIS), Hospital Universitario Virgen del Rocío, Departamento de Psiquiatría, Universidad de Sevilla, Sevilla, Spain; ⁷Biomedicine, Neuroscience and Advanced Diagnostic Department, Psychiatry Section, University of Palermo, Palermo, Italy; ⁸Department of Health Service and Population Research, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; ⁹Addiction and Mental Health Group (AIM), Department of Psychology, University of Bath, Bath, UK;

¹⁰National Addiction Centre, Institute of Psychiatry, King's College London, London, UK; ¹¹Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA; ¹²Psylyfe Group, Division of Psychiatry, University College London, London, UK; ¹³Department of Mental Health and Addiction Services, ASP Crotona, Crotona, Italy; ¹⁴Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement Sciences, University of Verona, Verona, Italy; ¹⁵Department of Medical and Surgical Science, Psychiatry Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy; ¹⁶Institut Mondor de recherche biomédicale, Creteil, France; ¹⁷Etablissement Public de Santé Maison Blanche, Paris, France; ¹⁸Department of Medicine and Surgery, University of Milano Bicocca, Monza, Italy; ¹⁹Department of Mental Health and Addiction Services, ASST Lecco, Lecco, Italy; ²⁰Department of Neurosciences, Mental Health and Sensory Organs, Suicide Prevention Center, Sant'Andrea Hospital, Sapienza University of Rome, Rome, Italy; ²¹Rivierduinen Institute for Mental Health Care, Leiden, The Netherlands; ²²Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, Maastricht, The Netherlands; ²³Early Psychosis Section, AmsterdamUMC, Academic Medical Centre, University of Amsterdam, Meibergdreef 5, 1105 AZ Amsterdam, The Netherlands; ²⁴Department of Preventive Medicine, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil; ²⁵Department of Psychiatry, Servicio de Psiquiatría Hospital "Virgen de la Luz", Cuenca, Spain; ²⁶Department of Psychiatry, Psychiatric Genetic Group, Instituto de Investigación Sanitaria de Santiago de Compostela, Complejo Hospitalario Universitario de Santiago de Compostela, Santiago, Spain; ²⁷Department of Medicine, Psychiatry Area, School of Medicine, Universidad de Oviedo, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Oviedo, Spain; ²⁸Department of Psychiatry, School of Medicine, Universidad de Valencia, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Valencia, Spain; ²⁹Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, University of Barcelona, Institut d'Investigacions Biomèdiques August Pi I Sunyer (IDIBAPS), Biomedical Research Networking Centre in Mental Health (CIBERSAM),

- Barcelona, Spain; ³⁰Department of Child and Adolescent Psychiatry, Institute of Psychiatry and Mental Health, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM, CIBERSAM, Madrid, Spain; ³¹Reader; PsyLife Group, Division of Psychiatry, University College London, London, UK; ³²Department of Psychiatry, University of Cambridge, Cambridge, UK; ³³CAMEO Early Intervention Service, Cambridgeshire & Peterborough NHS Foundation Trust, Cambridge, UK; ³⁴Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, UK; ³⁵Department Psychiatry, Brain Centre Rudolf Magnus, Utrecht University Medical Centre, Utrecht, The Netherlands; ³⁶Department of Psychiatry, Centre for PanorOmic Sciences, and State Key Laboratory of Brain and Cognitive Sciences, Li KaShing Faculty of Medicine, The University of Hong Kong, Hong Kong, China and ³⁷Research Foundation, National Institute for Health Research Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and the NIHR BRC at University College London, London, UK
- ## References
- Arseneault, L., Cannon, M., Poulton, R., Murray, R. M., Caspi, A., & Moffitt, T. E. (2002). Cannabis use in adolescence and risk for adult psychosis: Longitudinal prospective study. *BMJ*, *325*, 1212–1213. doi:10.1136/bmj.325.7374.1212.
- Barkus, E. J., Stirling, J., Hopkins, R. S., & Lewis, S. (2006). Cannabis-induced psychosis-like experiences are associated with high schizotypy. *Psychopathology*, *39*, 175–178. doi:10.1159/000092678.
- Bianconi, F., Bonomo, M., Marconi, A., Koliakou, A., Stilo, S. A., Iyegbe, C., ... Di Forti, M. (2016). Differences in cannabis-related experiences between patients with a first episode of psychosis and controls. *Psychological Medicine*, *46*, 995–1003. doi:10.1017/S0033291715002494.
- Buchan, B. J., Dennis, M. L., Tims, F. M., & Diamond, G. S. (2002). Cannabis use; consistency and validity of self-report, on-site urine testing and laboratory testing. *Addiction*, *97*, 98–108. doi:10.1046/j.1360-0443.97.s01.1.x.
- Dekker, N., Koeter, M., Van Den Brink, W., & Investigators, G. (2012). Craving for cannabis in patients with psychotic disorder, their non-affected siblings and healthy controls: Psychometric analysis of the obsessive compulsive drug use scale. *International Journal of Methods in Psychiatric Research*, *21*, 286–300. doi:10.1002/mpr.1362.
- Dekker, N., Linszen, D. H., & De Haan, L. (2009). Reasons for cannabis use and effects of cannabis use as reported by patients with psychotic disorders. *Psychopathology*, *42*, 350–360. doi:10.1159/000236906.
- Dekker, N., Smeerdijk, A. M., Wiers, R. W., Duits, J. H., van Gelder, G., Houben, K., ... de Haan, L. (2010). Implicit and explicit affective associations towards cannabis use in patients with recent-onset schizophrenia and healthy controls. *Psychological Medicine*, *40*, 1325–1336. doi:10.1017/S0033291709991814.
- Di Forti, M., Marconi, A., Carra, E., Fraietta, S., Trotta, A., Bonomo, M., ... Murray, R. M. (2015). Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis: A case-control study. *The Lancet Psychiatry*, *2*, 233–238. doi:10.1016/s2215-0366(14)00117-5.
- Di Forti, M., Morgan, C., Dazzan, P., Pariante, C., Mondelli, V., Marques, T. R., ... Murray, R. M. (2009). High-potency cannabis and the risk of psychosis. *The British Journal of Psychiatry*, *195*, 488–491. doi:10.1192/bjp.bp.109.064220.
- Di Forti, M., Quattrone, D., Freeman, T. P., Tripoli, G., Gayer-Anderson, C., Quigley, H., ... van der Ven, E. (2019). The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): A multicentre case-control study. *The Lancet Psychiatry*, *6*, 427–436. doi:10.1016/s2215-0366(19)30048-3.
- Dragt, S., Nieman, D. H., Becker, H. E., van de Fliert, R., Dingemans, P. M., de Haan, L., ... Linszen, D. H. (2010). Age of onset of cannabis use is associated with age of onset of high-risk symptoms for psychosis. *Canadian Journal of Psychiatry*, *55*, 165–171. doi:10.1177/070674371005500308.
- Etain, B., Lajnef, M., Bellivier, F., Henry, C., M'bailara, K., Kahn, J., ... Fisher, H. (2017). Revisiting the association between childhood trauma and psychosis in bipolar disorder: A quasi-dimensional path-analysis. *Journal of Psychiatric Research*, *84*, 73–79. doi:10.1016/j.jpsychires.2016.09.022.
- Ferdinand, R. F., Sondeijker, F., van der Ende, J., Selten, J. P., Huizink, A., & Verhulst, F. C. (2005). Cannabis use predicts future psychotic symptoms, and vice versa. *Addiction*, *100*, 612–618. doi:10.1111/j.1360-0443.2005.01070.x.
- Ferraro, L., La Cascia, C., La Barbera, D., Sanchez-Gutierrez, T., Tripoli, G., Seminerio, F., ... Quattrone, D. (2021). The relationship of symptom dimensions with premorbid adjustment and cognitive characteristics at first episode psychosis: Findings from the EU-GEI study. *Schizophrenia Research*, *236*, 69–79. doi:10.1016/j.schres.2021.08.008.
- Ferraro, L., La Cascia, C., Quattrone, D., Sideli, L., Matranga, D., Capuccio, V., ... Di Forti, M. (2020). Premorbid adjustment and IQ in patients with first-episode psychosis: A multisite case-control study of their relationship with cannabis use. *Schizophrenia Bulletin*, *46*, 517–529. doi:10.1093/schbul/sbz077.
- Ferraro, L., Russo, M., O'Connor, J., Wiffen, B. D., Falcone, M. A., Sideli, L., ... Di Forti, M. (2013). Cannabis users have higher premorbid IQ than other patients with first onset psychosis. *Schizophrenia Research*, *150*, 129–135. doi:10.1016/j.schres.2013.07.046.
- Freeman, T. P., Craft, S., Wilson, J., Stylianou, S., ElSohly, M., Di Forti, M., & Lynskey, M. T. (2021). Changes in delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) concentrations in cannabis over time: Systematic review and meta-analysis. *Addiction*, *116*, 1000–1010. doi:10.1111/add.15253.
- Freeman, T. P., Morgan, C. J., Hindocha, C., Schafer, G., Das, R. K., & Curran, H. V. (2014). Just say 'know': How do cannabinoid concentrations influence users' estimates of cannabis potency and the amount they roll in joints? *Addiction*, *109*, 1686–1694. doi:10.1111/add.12634.
- Gayer-Anderson, C., Jongsma, H. E., Di Forti, M., Quattrone, D., Velthorst, E., de Haan, L., ... Morgan, C. (2020). The European network of national schizophrenia networks studying gene-environment interactions (EU-GEI): Incidence and first-episode case-control programme. *Social Psychiatry and Psychiatric Epidemiology*, *55*, 645–657. doi:10.1007/s00127-020-01831-x.
- Gobbi, G., Atkin, T., Zytynski, T., Wang, S., Askari, S., Boruff, J., ... Mayo, N. (2019). Association of cannabis use in adolescence and risk of depression, anxiety, and suicidality in young adulthood: A systematic review and meta-analysis. *JAMA Psychiatry*, *76*, 426–434. doi:10.1001/jamapsychiatry.2018.4500.
- Green, B., Kavanagh, D. J., & Young, R. M. (2004). Reasons for cannabis use in men with and without psychosis. *Drug and Alcohol Review*, *23*, 445–453. doi:10.1080/09595230412331324563.
- Hjorthoj, C., Larsen, M. O., Starzer, M. S. K., & Nordentoft, M. (2021a). Annual incidence of cannabis-induced psychosis, other substance-induced psychoses and dually diagnosed schizophrenia and cannabis use disorder in Denmark from 1994 to 2016. *Psychological Medicine*, *51*, 617–622. doi:10.1017/S0033291719003532.
- Hjorthoj, C., Posselt, C. M., & Nordentoft, M. (2021b). Development over time of the population-attributable risk fraction for cannabis use disorder in schizophrenia in Denmark. *JAMA Psychiatry*, *78*, 1013–1019. doi:10.1001/jamapsychiatry.2021.1471.
- Jongsma, H. E., Gayer-Anderson, C., Lasalvia, A., Quattrone, D., Mule, A., & Szoke, A., ... European Network of National Schizophrenia Networks Studying Gene-Environment Interactions Work Package, G. (2018). Treated incidence of psychotic disorders in the multinational EU-GEI study. *JAMA Psychiatry*, *75*, 36–46. doi:10.1001/jamapsychiatry.2017.3554.
- Khantzian, E. J. (1985). The self-medication hypothesis of addictive disorders: Focus on heroin and cocaine dependence. *The American Journal of Psychiatry*, *142*, 1259–1264. doi:10.1176/ajp.142.11.1259.
- Koliakou, A., Joseph, C., Ismail, K., Atakan, Z., & Murray, R. M. (2011). Why do patients with psychosis use cannabis and are they ready to change their use? *International Journal of Developmental Neuroscience*, *29*, 335–346. doi:10.1016/j.ijdevneu.2010.11.006.
- Korver, N., Nieman, D. H., Becker, H. E., van de Fliert, R., Dingemans, P. H., de Haan, L., ... Linszen, D. H. (2010). Symptomatology and neuropsychological functioning in cannabis using subjects at ultra-high risk for developing psychosis and healthy controls. *Australian & New Zealand Journal of Psychiatry*, *44*, 230–236. doi:10.3109/00048670903487118.
- Kwok, O.-M., Cheung, M. W., Jak, S., Ryu, E., & Wu, J.-Y. (2018). Recent advancements in structural equation modeling (sem): From both methodological and application perspectives. *Frontiers in Psychology*, *9*, 1936. doi:10.3389/fpsyg.2018.01936.
- Leung, J., Chan, G., Stjepanovic, D., Chung, J. Y. C., Hall, W., & Hammond, D. (2022). Prevalence and self-reported reasons of cannabis use for medical

- purposes in USA and Canada. *Psychopharmacology (Berl)*, 239, 1509–1519. doi:10.1007/s00213-021-06047-8.
- Mallett, R., Leff, J., Bhugra, D., Pang, D., & Zhao, J. H. (2002). Social environment, ethnicity and schizophrenia. A case-control study. *Social Psychiatry and Psychiatric Epidemiology*, 37, 329–335. doi:10.1007/s00127-002-0557-4.
- Marconi, A., Di Forti, M., Lewis, C. M., Murray, R. M., & Vassos, E. (2016). Meta-analysis of the association between the level of cannabis use and risk of psychosis. *Schizophrenia Bulletin*, 42, 1262–1269. doi:10.1093/schbul/sbw003.
- Martin, A. J. (2011). Prescriptive statements and educational practice: What can structural equation modeling (SEM) offer? *Educational Psychology Review*, 23, 235–244. doi:10.1007/s10648-011-9160-0.
- Maxwell, M. E. (1992). *Family interview for genetic studies (FIGS): A manual for FIGS*. Bethesda, MD: Clinical Neurogenetics Branch, Intramural Research Program, National Institute of Mental Health.
- McGuffin, P., Farmer, A. E., & Harvey, I. (1991). A polydiagnostic application of operational criteria in studies of psychotic illness: Development and reliability of the OPCRIT system. *Archives of General Psychiatry*, 48, 764–770. doi:10.1001/archpsyc.1991.01810320088015.
- Murray, R. M., Englund, A., Abi-Dargham, A., Lewis, D. A., Di Forti, M., Davies, C., ... D'Souza, D. C. (2017). Cannabis-associated psychosis: Neural substrate and clinical impact. *Neuropharmacology*, 124, 89–104. doi:10.1016/j.neuropharm.2017.06.018.
- Mustonen, A., Niemelä, S., Nordström, T., Murray, G. K., Mäki, P., Jääskeläinen, E., & Miettunen, J. (2018). Adolescent cannabis use, baseline prodromal symptoms and the risk of psychosis. *The British Journal of Psychiatry*, 212, 227–233. doi:10.1192/bjp.2017.52.
- Pérez, L. G., Santacana, A. N., Baquero, D. B., & Pérez-Solà, V. (2014). Reasons and subjective effects of cannabis use among people with psychotic disorders: A systematic review. *Acta Espanolas de Psiquiatria*, 42, 83–90. Retrieved from <https://pubmed.ncbi.nlm.nih.gov/24715366/>.
- Peters, B. D., de Koning, P., Dingemans, P. H., Becker, H. E., Linszen, D. H., & de Haan, L. (2009). Subjective effects of cannabis before the first psychotic episode. *Australian and New Zealand Journal of Psychiatry*, 43, 1115–1162. doi:10.3109/00048670903179095.
- Radhakrishnan, R., Pries, L. K., Erzin, G., Ten Have, M., de Graaf, R., van Dorsselaer, S., ... Guloksuz, S. (2022). Bidirectional relationships between cannabis use, anxiety and depressive symptoms in the mediation of the association with psychotic experience: Further support for an affective pathway to psychosis. *Psychological Medicine*, 1–7. doi:10.1017/S0033291722002756.
- Radhakrishnan, R., Wilkinson, S. T., & D'Souza, D. C. (2014). Gone to Pot - A review of the association between cannabis and psychosis. *Frontiers in Psychiatry*, 5, 54. doi:10.3389/fpsy.2014.00054.
- Stage, F. K., Carter, H. C., & Nora, A. (2004). Path analysis: An introduction and analysis of a decade of research. *The Journal of Educational Research*, 98, 5–13. doi:10.3200/joer.98.1.5-13.
- Tripoli, G., Quattrone, D., Ferraro, L., Gayer-Anderson, C., Rodriguez, V., La Cascia, C., ... Di Forti, M. (2021). Jumping to conclusions, general intelligence, and psychosis liability: Findings from the multi-centre EU-GEI case-control study. *Psychological Medicine*, 51, 623–633. doi:10.1017/S003329171900357X.
- van Os, J., Pries, L. K., Ten Have, M., de Graaf, R., van Dorsselaer, S., Bak, M., ... Guloksuz, S. (2021). Schizophrenia and the environment: Within-person analyses may be required to yield evidence of unconfounded and causal association-The example of cannabis and psychosis. *Schizophrenia Bulletin*, 47, 594–603. doi:10.1093/schbul/sbab019.
- Williams, J., Farmer, A. E., Ackenheil, M., Kaufmann, C. A., & McGuffin, P. (1996). A multicentre inter-rater reliability study using the OPCRIT computerized diagnostic system. *Psychological Medicine*, 26, 775–783. doi:10.1017/s003329170003779x.