

Research article

Open Access

## Genetic transformation of *Vitis vinifera* via organogenesis

Bruno Mezzetti\*<sup>1</sup>, Tiziana Pandolfini<sup>2</sup>, Oriano Navacchi<sup>3</sup> and Lucia Landi<sup>1</sup>

Address: <sup>1</sup>Dipartimento Biotecnologie Agrarie e Ambientali, University of Ancona, Via Breccie Bianche 60100 – Ancona Italy, <sup>2</sup>Dipartimento Scientifico Tecnologico, University of Verona, Strada Le Grazie, 37134 Verona Italy and <sup>3</sup>Vitroplant, Via Loreto – 47023 Cesena Italy

E-mail: Bruno Mezzetti\* - [bruno@unian.it](mailto:bruno@unian.it); Tiziana Pandolfini - [tiziana.pandolfini@univr.it](mailto:tiziana.pandolfini@univr.it); Oriano Navacchi - [info@vitroplant.it](mailto:info@vitroplant.it); Lucia Landi - [ebpatveg@mta01.unian.it](mailto:ebpatveg@mta01.unian.it)

\*Corresponding author

Published: 27 September 2002

Received: 7 May 2002

*BMC Biotechnology* 2002, **2**:18

Accepted: 27 September 2002

This article is available from: <http://www.biomedcentral.com/1472-6750/2/18>

© 2002 Mezzetti et al; licensee BioMed Central Ltd. This article is published in Open Access: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

**Keywords:** *Vitis vinifera*, meristem regeneration, genetic transformation, gene expression, DefH9-iaaM gene

### Abstract

**Background:** Efficient transformation and regeneration methods are a priority for successful application of genetic engineering to vegetative propagated plants such as grape. The current methods for the production of transgenic grape plants are based on *Agrobacterium*-mediated transformation followed by regeneration from embryogenic callus. However, grape embryogenic calli are laborious to establish and the phenotype of the regenerated plants can be altered.

**Results:** Transgenic grape plants (*V. vinifera*, table-grape cultivars Silcora and Thompson Seedless) were produced using a method based on regeneration via organogenesis. *In vitro* proliferating shoots were cultured in the presence of increasing concentrations of N<sup>6</sup>-benzyl adenine. The apical dome of the shoot was removed at each transplantation which, after three months, produced meristematic bulk tissue characterized by a strong capacity to differentiate adventitious shoots. Slices prepared from the meristematic bulk were used for *Agrobacterium*-mediated transformation of grape plants with the gene *DefH9-iaaM*. After rooting on kanamycin containing media and greenhouse acclimatization, transgenic plants were transferred to the field. At the end of the first year of field cultivation, *DefH9-iaaM* grape plants were phenotypically homogeneous and did not show any morphological alterations in vegetative growth. The expression of *DefH9-iaaM* gene was detected in transgenic flower buds of both cultivars.

**Conclusions:** The phenotypic homogeneity of the regenerated plants highlights the validity of this method for both propagation and genetic transformation of table grape cultivars. Expression of the *DefH9-iaaM* gene takes place in young flower buds of transgenic plants from both grape cultivars.

### Background

The use of genetic engineering for plant improvement permits the introduction of useful agronomic traits without altering the features of the cultivar, necessitating the development of *in vitro* systems for the genetic transformation and plant regeneration. To date, the regeneration of

grape plants has been obtained by both organogenesis and embryogenesis. Shoot regeneration from fragmented shoot apices has been successfully applied to several grape species and hybrids [1]. Moreover, adventitious shoots have been regenerated from either leaf blades or petioles of *V. rupestris*, *V. vinifera* and selected grape hybrids [2].

Embryogenic cultures have also been established for some grape species, e.g. *Vitis longii* Prince [3], *V. rupestris* Scheele [4,5], *V. x lambruscana* [6] and *V. vinifera* varieties [7,4,8] and rootstock hybrids [9].

Regeneration from somatic embryos has been used for genetic transformation of the major grape cultivars for production of both wine [10] and table grapes [11,12]. Transgenic grape can be generated by transformation of embryogenic calli obtained from different tissues, including zygotic embryos [11], leaves [13], ovaries [6] and anther filaments. The latter is the most widely used tissue for these purposes [3,10,12]. The initiation and proliferation of embryogenic callus from anther filaments is, however, laborious and cultivar-dependent [14]. Furthermore, regeneration by somatic embryogenesis is limited by the availability of immature flowers for the initial explant and strongly affects the phenotype of grapevine plants [6,12,15].

Herein, an alternative method useful for both propagation and *Agrobacterium*-mediated genetic transformation of table grape (*V. vinifera*) is described. The method is based on the formation of meristematic bulk (MB) tissue with a high regenerative capacity, using adventitious shoots as a starting material. This procedure has been used to introduce the *DefH9-iaaM* gene [17] into the genome of two table grape cultivars (Silcora and Thompson Seedless). The aforementioned gene is composed of the ovule-specific regulatory regions from *DefH9* of *Antirrhinum majus* and the *iaaM* coding region from *Pseudomonas savastanoi*. The *DefH9-iaaM* gene has conferred parthenocarpic fruit development to several horticultural species improving fruit production [18–22]. The transformed grape plants show normal vegetative growth and express the *DefH9-iaaM* gene in young flower buds.

## Results

### Initiation and maintenance of meristematic bulk

The method has been used to genetically engineer the table grape varieties Silcora and Thompson Seedless. The method combines two *in vitro* processes, namely the production of meristematic tissue with a high competence for plant regeneration and the genetic transformation of this tissue, followed by selection of transformed regenerants (Fig. 1).

The first process generates the meristematic bulk tissue, a cellular aggregate with an elevated regenerative capacity. The protocol used to produce meristematic bulk tissue consists in four subculturing phases starting from *in vitro* proliferating shoots. In the first phase, the apical dome of the shoots are eliminated and the basal cluster maintained for 30 days in culture medium (IM) supplemented with 4.4  $\mu\text{M}$  BA. In the second phase, after a further drastic

dissection of the apical dome, the shoots are transplanted to IM medium containing 8.8  $\mu\text{M}$  BA and cultured for 30 days. In the third and fourth phases, the mechanical dissection of the apical dome is repeated after 60 and 90 days of culture, respectively, and the concentration of BA in the medium is further increased to 13.2  $\mu\text{M}$ .

The meristematic bulk can be cultured for long periods of time on MM medium (13.2  $\mu\text{M}$  BA) with the continuous formation of new adventitious buds that are clearly visible at the surface of the aggregate (Fig. 2A). The internal part of the meristematic bulk tissue shows hypertrophy of parenchyma cells with highly vascularized bands (Fig. 2B), including many initiation nodules (Fig. 2C) from which adventitious buds originate (Fig. 2D).


This meristematic bulk tissue can be easily propagated: the tissue is fragmented and each fragment generates a new meristematic bulk tissue after 3–4 weeks of culture. Moreover, by cutting the meristematic bulk tissue in small slices (1  $\text{cm}^2$ , 2 mm thick), tissue layers (Fig. 3A) that produce high numbers of regenerated adventitious shoots (about 20 per slice within four weeks of culture on MM medium) are obtained (Fig. 3B). These meristematic slices are suitable as a starting material for both vegetative plant propagation and genetic transformation.

### Genetic Transformation and Selection

Grape plants of the cultivars Silcora and Thompson Seedless were transformed with the *DefH9-iaaM* gene. Meristematic slices of the two cultivars were inoculated with *agrobacteria* harboring the *DefH9-iaaM* construct. After 30 days on medium containing 25  $\text{mg l}^{-1}$  kanamycin, a high number of regenerants were obtained. When transferred to medium containing higher concentrations of the antibiotic (50  $\text{mg l}^{-1}$ ), many of the regenerants became yellowish, and were discarded (60 days). The selection of stable regenerants was obtained after an additional 30-day incubation on medium with an even higher concentration of kanamycin (75  $\text{mg l}^{-1}$ ). During *in vitro* selection, only the transformed cells maintain a high regenerative capacity, while the other parts of the meristematic tissue become progressively necrotic (Fig. 4). Three to five regenerants were isolated per meristematic slice and transferred to rooting medium containing 75  $\text{mg l}^{-1}$  kanamycin. About seven months following the initiation of MB tissue culture, 50 putative transgenic lines of the two grape cultivars were isolated.

### Transgenic state and gene expression


The transgenic state of 10 *in vitro* rooted plants of the cultivar Silcora and five plants of the cultivar Thompson Seedless was determined by Southern blot analysis. Eight out of ten plants of the cultivar Silcora were transgenic for the *DefH9-iaaM* gene. The eight transformants arose from



**Figure 1**  
Grape micropropagation and genetic transformation: schematic representation of the *in vitro* processes. BA, benzyl adenine; MB, meristematic bulk.

three independent events: #6, #29, and #35, had 2, 3, and 1 copies of the transgene, respectively (Fig. 5A, lanes 3, 5, and 6). Of the five Thompson Seedless plants analyzed, three were transgenic for *DefH9-iaaM*. All the transgenic plants were recovered from a unique transformation event (line #4) and had a single copy of the transgene (Fig. 5A, lane 1).

The expression of the *DefH9-iaaM* gene in transgenic flower buds was analyzed by RT-PCR (Fig. 6). An amplicon of 266 bp corresponding to a fragment of the spliced *DefH9-iaaM* transcript was detected when mRNA extracted from flower buds of *DefH9-iaaM* transgenic Thompson Seedless plants (line #4, Fig 6, lane 2) and Silcora plants (lines #6 and #35, Fig. 6 lanes 4 and 5, respectively) was used as template for RT-PCR analysis. No amplicon was detected using mRNA extracted from flower buds of untrans-



**Figure 2**  
Meristematic bulk tissue (MB) generated from *in vitro*-proliferated grape shoots: (A) standard proliferation stage; (B) Internal section ( $\times 40$ ) of the MB showing hyper-trophic parenchymatous cells with highly vascularized bands; many initiation nodules ( $\times 40$ ) are also visible (C). The nodules generate adventitious buds ( $\times 5$ ) (D).


formed Silcora and Thompson Seedless plants as a template (Fig 6, lanes 1 and 3). Thus, in both cultivars the inserted transgene is expressed in the proper organ.

#### **Phenotype of transgenic plants regenerated from meristematic bulks**

The transgenic plants of the two table grape varieties were proliferated *in vitro* and after acclimatization in the greenhouse, they were then transferred to the field as self-rooted plants. Forty plants for each independent line were grown (Silcora lines #6 and #35 and Thompson Seedless line #4). During field development, the plants showed normal vegetative growth and appeared healthy. Transgenic and control lines of both Thompson Seedless and Silcora showed a high similarity in leaf morphology (Fig. 7 and 8). The similarity is already evident in the first expanded leaves (Fig 7A Thompson Seedless left: control,

right: line #4; Fig 7B Silcora centre: control, left and right: lines #6 and #35, respectively). The comparison of adult leaves (from the 6<sup>th</sup> to 12<sup>th</sup> position on the branch) of transgenic and control lines from either Thompson Seedless (Fig. 8 panel A: control; panel B: line #4) or Silcora (Fig. 8 panel C: control; panel D: line #6; panel E: line #35) confirmed their morphological resemblance.

Transgenic Thompson Seedless plants show the main features reported in its ampelographic description [23]. The only and rather trivial difference observed in adult leaves of the transgenic line was the presence of slightly deeper lower-lateral sinuses in comparison to control (Fig. 8B). It is worth noticing that the leaves of both cultivars had neither lobes nor red-vein, alterations observed in plants regenerated *in vitro* by Franks et al. [12]. The *DefH9-iaaM* grape plants are currently under agronomic (i.e. field trial)



**Figure 3**  
Grape regeneration via organogenesis: (A) slices (approx. 1 cm<sup>2</sup>, 2 mm thick) prepared from the MB and used for propagation or genetic transformation; (B) shoot regeneration obtained after 30 days of culture ( $\times 5$ ).



**Figure 4**  
Regenerating transgenic lines on the selection medium supplemented with 50 mg/l kanamycin 60 days after infection.

evaluation for fruit development and total fruit production.

### Discussion


Transgenic grape plants are typically regenerated from somatic embryos derived from either zygotic embryos or leaves [11,13], or more recently, from anther filaments [10,12]. In our method, transgenic grape plants are regenerated via organogenesis; a meristematic tissue with high regenerative capacity is established from *in vitro* standard proliferating shoots and used for both propagation and *A. tumefaciens*-mediated transformation.

Combined with the removal of the shoot apices, the progressive increase of cytokinin content in the media abolishes the shoot apical dominance and also promotes basal proliferation, leading to the formation of meristematic bulk tissue. This is a large aggregate of meristematic tissue, which includes parenchymatous as well as primary and secondary meristematic cells. The tissue layers prepared by slicing the meristematic bulk show a high meristematic/regenerative competence and within four weeks, produce a high number of regenerated adventitious shoots (about 20 per slice).

This meristematic tissue is produced in a relatively short period of time (90 days) and is valuable for both vegetative plant propagation and genetic transformation. For the success of *Agrobacterium* mediated transformation, the slicing of the meristematic bulk is a critical aspect, producing tissue composed of cells having a high regenerative capacity and a large number of damaged cells. This latter population of cells is known to produce substances that activate the mechanism of genetic transfer from *Agrobacterium* [24].

The main advantage of this transformation method is represented by the shoot regeneration efficiency, which is higher than that achieved by organogenesis using lateral axillary shoots or leaf. In fact, an average of 3–5 regenerants per shoot and 0.1–0.5 regenerant per leaf are produced [25]. On the other hand, a selection strategy based on the progressive increase of the concentration of kanamycin was designed to limit the number of escapes or chimerical events. For both cultivars, untransformed kanamycin resistant plants represented an average of 20% of all plants analyzed.

High concentrations of kanamycin cause oxidative damage to non-resistant tissues and lead to progressive necrosis. It has been reported that the presence of necrotic areas in transformed grape embryogenic tissue may affect the regeneration efficiency [14]. In our system, although the highest concentration of kanamycin (75 mg l<sup>-1</sup>) induced an extensive browning and necrotization of the meristem-



**Figure 5**

Southern blot analysis of grape plants transgenic for *DefH9-iaaM* gene. (A) Genomic DNA (digested with *HindIII*) from control untransformed Silcora plants (lanes 4, 7), three independent transgenic Silcora lines (plants #6, #29, #35, lanes 3, 5, 6, respectively), control untransformed Thompson Seedless plants (lane 2) and transgenic Thompson Seedless line (plant #4, lane 1). (B) Schematic drawings of the constructs used for transformation of Silcora (right) and Thompson Seedless (left) plants are reported. The probes are indicated with grey boxes. Only restriction sites relevant for Southern analysis are indicated. LB, left border; R, right border.


atic slices, the regeneration capacity of the transformed cells was unaltered.

The production of transgenic grape plantlets requires four months of culture starting from meristematic bulk; seven months are needed if *in vitro* proliferating shoots are used as the starting material. Grape transformation methods based on the induction and selection of somatic embryos are comparable in length [10]. However, they appear to be more laborious with respect to the *in vitro* manipulations required for regeneration (establishment of embryogenic callus and embryos selection, maturation and regeneration) [6]. They also present a disadvantage regarding the

availability of the starting material (in particular anthers filaments).

At the end of the first year of field cultivation, the transgenic grape plants were phenotypically homogeneous and highly similar to the control plants. Moreover they did not show the alterations observed in grape plants obtained with different methods of regeneration, such as stronger anthocyanin pigmentation, reduced vigor, leaf lobation, and albinism [12,15,16].

It is worth noting that the transgene was expressed in the flower buds of *DefH9-iaaM* plants of both grape cultivars. Thus to date, the *DefH9-iaaM* gene has been shown to be




**Figure 6**  
RT-PCR analysis of flower buds from transgenic grape plants. Analysis was performed with single strand cDNA synthesized from mRNA extracted from young flower buds of Silcora control and transgenic plants #6 and #35 (lanes 3, 4 and 5, respectively) and Thompson Seedless control and transgenic plants (lanes 1 and 2, control and #4, respectively). The amplification product of 266 bp corresponds to the 5' end of the spliced *DefH9-iaaM* mRNA.

properly expressed in three different plant families: *Solanaceae* [17,18], *Rosaceae* [21], *Vitaceae* (present manuscript).

## Conclusions

The present report describes a new method for table grape (*Vitis vinifera*) genetic transformation. This method of transformation differs from previously described techniques with respect to regeneration (based on organogenesis rather than embryogenesis) and it is characterized by a higher regeneration efficiency, achieved through simple *in vitro* manipulations. Moreover, the phenotypic stability and homogeneity of the regenerated plants during the vegetative growth, highlights the validity of this method for both propagation and genetic transformation of Thompson Seedless and Silcora table grape cultivars. It is likely that the method can be applied to other grape cultivars, for example cultivars used in the wine industry. The transgenic plants express the transgene in the proper organ. Agronomic evaluation of the productive performance of *DefH9-iaaM* transgenic grapes will permit to evaluate the utility of this molecular tool in grape scion improvement.



**Figure 7**  
First expanded leaves of transgenic and control lines of Thompson Seedless (A. left: control; right: line #4) and Silcora (B left: line #6; centre: control; right: line #35).


## Methods

### Plant material

*In vitro* proliferating cultures of *Vitis vinifera*, table grape varieties Silcora (I.G. 235023) and Thompson Seedless, were initiated from shoot-tips and subcultured monthly on propagation medium (PR). This medium is composed of MS basal medium [26] supplemented with 4.4  $\mu\text{M}$  benzyl adenine (BA), 3 % sucrose, and 0.7% commercial agar (pH 5.6). The cultures were kept at  $25 \pm 1^\circ\text{C}$  under a photoperiod of 16 h light ( $70 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  photosynthetic photon flux) provided by warm white fluorescent tubes.

### Initiation and maintenance of meristematic bulks

Proliferating shoots were subjected to chemical and mechanical treatments to induce the formation of meristematic bulks characterized by a strong capacity to differentiate adventitious shoots. The medium used for the initiation (IM) of the meristematic bulk (MB) contained  $\text{KNO}_3$  (1050  $\text{mg l}^{-1}$ ),  $\text{NH}_4\text{NO}_3$  (400  $\text{mg l}^{-1}$ ),  $\text{KH}_2\text{PO}_4$  (200  $\text{mg l}^{-1}$ ),  $\text{MgSO}_4 \cdot 7 \cdot \text{H}_2\text{O}$  (400  $\text{mg l}^{-1}$ ),  $\text{CaNO}_3$  (750  $\text{mg l}^{-1}$ ),  $\text{NaH}_2\text{PO}_4$  (200  $\text{mg l}^{-1}$ ), microele-



**Figure 8**  
Adult leaves (collected from the 6<sup>th</sup> to 12<sup>th</sup> position on the branch) of transgenic and control lines of Thompson Seedless (A: control; B: line # 4) and Silcora (C: control; D: Line # 6; E: Line # 35).

ments and vitamins by MS [26], 3% sucrose, 0.7% commercial agar, and 0.05  $\mu\text{M}$  NAA. IM medium was supplemented with 4.4  $\mu\text{M}$  BA for the first 30 day subculture, after which the BA concentration was doubled (8.8  $\mu\text{M}$ ) for the second 30 day subculture. MB was maintained on IM medium supplemented with 13.2  $\mu\text{M}$  BA (MM medium) and subcultured every 4 weeks.

The mechanical treatment consisted in the elimination of the apical dome of the initial proliferating shoots. This procedure was repeated, at each transplantation, until the MB was obtained. The inner part of the MB was cut in thin slices (1  $\text{cm}^2$ , 2 mm thick), which were transferred to fresh MM medium and used for both micropropagation and genetic transformation.

#### Histological study

For histological analysis, the meristematic bulk tissue was cut using a cryo-microtome. The resulting 20  $\mu\text{m}$ -thick slices were stained with the DAPI fluorescent staining [27]. The samples were analyzed using a Nikon microscope equipped with an epifluorescent light source.

#### Genetic transformation

Transformation experiments were carried out using *A. tumefaciens* strain C58 GV3101 [28] harbouring the parthenocarpic gene *DefH9-iaaM* [17] cloned in the pPCV002 binary vector [28]. The T-DNA of the vector contains the *nptII* coding region, conferring kanamycin resistance as selectable marker. Bacteria were cultured in the YEB medium (1 g  $\text{l}^{-1}$  yeast extract, 5 g  $\text{l}^{-1}$  beef extract, 5 g  $\text{l}^{-1}$  peptone, 5 g  $\text{l}^{-1}$  sucrose and 2 mM  $\text{MgSO}_4$ , pH 7.8) containing 50 mg  $\text{l}^{-1}$  kanamycin, 100 mg  $\text{l}^{-1}$  rifampicin, and 100 mg  $\text{l}^{-1}$  streptomycin. Bacterial cultures ( $A_{420}$  0.5) were grown overnight (29°C – 150 rpm), centrifuged, and resuspended in MS salts supplemented with 2% sucrose, 1 mM acetosyringone, 1 mM proline (pH 5.2) for 5 hrs at 28°C. Slices (1  $\text{cm}^2$ , 2 mm thick) obtained from the MB were dipped in the bacterial suspension for 15 min. After infection, the slices were blotted with sterile filter paper and placed on MM medium. After 48 hrs at 28°C, the explants were transferred to the same medium enriched with 25 mg  $\text{l}^{-1}$  kanamycin and 200 mg  $\text{l}^{-1}$  cefotaxime and maintained for 30 days. The explants were then transferred and subcultured on a monthly basis on the same medium with increasing concentrations of kanamycin (50 and 75 mg  $\text{l}^{-1}$ ). Detectable regenerants were isolated and transferred to rooting medium containing macroelements by Quoirin and Lepoivre [28]; microelements and vitamins by MS [26]; 4.9  $\mu\text{M}$  IBA; 5.7  $\mu\text{M}$  IAA; 3% sucrose, and 0.7% commercial agar, supplemented with 75 mg  $\text{l}^{-1}$  of kanamycin.

The *in vitro* rooted transgenic clones of Silcora and Thompson Seedless were grown in a greenhouse for accli-

matization (30 days) and weaning (60 days) and then transferred to the experimental field. Untransformed plants produced by standard micropropagation techniques were used as controls in field experiments.

#### Southern analysis

Genomic DNA was extracted from 1 g of frozen leaves or shoot apices using the Nucleon PhytoPure system (Amersham Pharmacia) according to the manufacturer's instructions. Ten  $\mu\text{g}$  of DNA from transgenic plants was digested with 70 Units of *HindIII*. The DNA was subjected to electrophoresis on a 0.7% agarose gel at 4.5  $\text{V cm}^{-1}$  and transferred to a nylon membrane (Hybond N, Amersham). The membrane was hybridized with 100 ng of fluorescein-labeled probe prepared using the Random Prime Labeling Module kit from Amersham. The probe was either a 600 bp-long DNA fragment of the *iaaM* coding region (Silcora plants) or a 2130 bp-long DNA fragment of the *DefH9* promoter (Thompson plants). Detection was performed with anti-fluorescein AP conjugate (Amersham) and the chemiluminescent alkaline phosphatase CDP-Star substrate (Amersham) according to the manufacturer's instructions. The membranes were exposed for 1 h using Kodak XAR-5 film.

#### RT-PCR analysis

Flower buds (0.5 cm long) were frozen in liquid nitrogen and total RNA was extracted by using the NucleonPhytopure system (Amersham). The system was slightly modified by the addition of Polyclar AT (95 mg  $\text{g}^{-1}$  of fresh tissue) and  $\text{Na}_2\text{S}_2\text{O}_5$  (0.4 %) to the homogenization buffer. The RNA was recovered by LiCl precipitation. Poly(A<sup>+</sup>)RNA was isolated from total RNA using oligo d(T) Dynabeads (Dyna) following the manufacturer's protocol. The amount of mRNA extracted was determined spectrophotometrically. Messenger RNA (1  $\mu\text{g}$ ) was used as a template in the reverse transcriptase reaction primed with oligodT<sub>20</sub>.

First strand cDNA (30–50 ng) was amplified with the forward primer 5'-CTTTGGAAGCTCGTGTGAGCTCTCA-3' (corresponding to the *DefH9* ULR region +89 +113, with +1 the transcription initiation nucleotide). The 3' primer was 5'-ACTATCGCTACCCGAGGGGTGGGC 3', complementary to the *iaaM* coding region from +131 to +108 bp. The resulting amplicon (266 bp), corresponding to a fragment of the spliced *DefH9-iaaM* mRNA, was characterized by sequence analysis.

#### Abbreviations

BA: N<sup>6</sup>-benzyl adenine; IBA: indole-3-butyric acid; MB: meristematic bulk; MS: Murashige and Skoog; NAA: 1-naphthaleneacetic acid.

## Authors' Contributions

BM carried out the histological and phenotypic analysis, participated in the genetic transformation. TP carried out the molecular analysis. ON carried out the "in vitro" regeneration, proliferation and selection. LL carried out the genetic transformation and participated in the histological study. All authors read and approved the final manuscript.

## Acknowledgements

This work was in part financed by the program "Biotecnologie II" of the CNR. The authors thank Angelo Spena for helpful advice.

## References

- Barlass M, Skene KGM: **In vitro propagation of grapevine (*Vitis vinifera* L.) from fragmented shoot apices.** *Vitis* 1978, **17**:335-340
- Stamp JA, Meredith CP: **Somatic embryogenesis from leaves and anthers of grapevine.** *Scientia Horticulturae* 1988, **35**:235-250
- Mullins MG, Rajasekaran K: **Plantlets from cultured anthers of *Vitis* species and hybrids.** *Proc Third Int Symp Grape Breeding, Davis* 1980, 111-119
- Stamp JA, Colby SM, Meredith CP: **Direct shoot organogenesis and plant regeneration from leaves of grape (*Vitis* spp.).** *Plant Cell Tissue Organ Culture* 1990, **22**:127-133
- Martinelli L, Bragagna P, Poletti V, Scienza A: **Somatic embryogenesis from leaf- and petiole-derived callus of *Vitis rupestris*.** *Plant Cell Report* 1993, **12**:207-210
- Motoike SY, Skirvin RM, Norton MA, Otterbacher AG: **Somatic embryogenesis and long term maintenance of embryogenic lines from fox grape.** *Plant Cell Tissue Organ Culture* 2001, **66**:121-131
- Mauro MC, Nef C, Fallot J: **Stimulation of somatic embryogenesis and plant regeneration from anther culture of *Vitis vinifera* cv. Cabernet-Sauvignon.** *Plant Cell Report* 1986, **5**:377-380
- Matsuta N, Hirabayashi T: **Embryogenic cell lines from somatic embryos of grape (*Vitis vinifera* L.).** *Plant Cell Report* 1989, **7**:684-687
- Coutos-Thevenot P, Goebel-Tourand I, Mauro MC, Jouanneau JP, Boulay M, Deloire A, Guern J: **Somatic embryogenesis from grapevine cells: I. Improvement of embryo development by changes in culture conditions.** *Plant Cell Tissue Organ Culture* 1992, **29**:125-133
- Iocco P, Franks T, Thomas MR: **Genetic transformation of major wine grape cultivars of *Vitis vinifera* L.** *Transgenic Research* 2001, **10**(2):105-112
- Scorza R, Cordts JM, Ramming DW, Emershad RL: **Transformation of grape (*Vitis vinifera* L.) zygotic-derived somatic embryos and regeneration of transgenic plants.** *Plant Cell Reports* 1995, **14**(9):589-592
- Franks T, Gang He D, Thomas MR: **Regeneration of transgenic shape *Vitis vinifera* L. Sultana plants: genotypic and phenotypic analysis.** *Molecular Breeding* 1998, **4**:321-333
- Scorza R, Cordts JM, Gray DJ, Gonsalves D, Emershad RL, Ramming DW: **Producing transgenic 'Thompson Seedless' grape (*Vitis vinifera* L.) plants.** *Journal of the American Society for Horticultural Science* 1996, **121**(4):616-619
- Perl A, Lotan O, Abu-Abied M, Holland D: **Establishment of an *Agrobacterium*-mediated transformation system for grape (*Vitis vinifera* L.): the role of antioxidants during grape-*Agrobacterium* interactions.** *Nature Biotechnology* 1996, **14**(11):1521
- Rajasekaran K, Mullins MG: **Embryos and plantlets from cultured anthers of hybrid grapevines.** *Journal Experimental Botany* 1979, **30**:399-407
- Rajasekaran K, Mullins MG: **The origin of embryos and plantlets from cultured anther of hybrid grapevines.** *Journal Enology and Viticulture* 1983, **34**:108-113
- Rotino GL, Perri E, Zottini M, Sommer H, Spena A: **Genetic engineering of parthenocarpic plants.** *Nature Biotechnology* 1997, **15**:1398-2001
- Ficcadenti N, Sestili S, Pandolfini T, Cirillo C, Rotino GL, Spena A: **Genetic engineering of parthenocarpic fruit development in tomato.** *Molecular Breeding* 1999, **5**:463-470
- Donzella G, Spena A, Rotino GL: **Transgenic parthenocarpic eggplants: superior germoplasm for increased winter production.** *Molecular Breeding* 2000, **6**:79-86
- Pandolfini T, Rotino GL, Camerini S, Defez R, Spena A: **Optimization of transgene action at the post-transcriptional level: high quality parthenocarpic fruits in industrial tomatoes.** *BMC Biotechnology* 2002, **2**:1 [http://www.biomedcentral.com/1472-6750/2/1]
- Mezzetti B, Landi L, Scortichini L, Rebori A, Spena A, Pandolfini T: **Genetic engineering of parthenocarpic fruit development in strawberry.** *Acta Horticulturae* 2002, **567**:101-104
- Acciarri N, Restaino F, Vitelli G, Perrone D, Zottini M, Pandolfini T, Spena A, Rotino GL: **Improved fruit productivity under both greenhouse and open field cultivation.** *BMC Biotechnology* 2002, **2**:4 [http://www.biomedcentral.com/1472-6750/2/4]
- Galet P, Morton LT: **A practical Ampelography: Grapevine Identification.** Cornell University Press 1979
- Nester E, Yong-Woog L, Shouguang J, Woong Seop S: **The sensing of plant signal molecules by *Agrobacterium*.** In: *Molecular aspects of pathogenicity and resistance: requirement for signal trasduction* (Edited by: Mills D, Kumoh H, Keen NT and Mayama S) APS Press 1996, 95-103
- Navacchi O, Mezzetti B, Zuccherelli G, Spena A: **Metodo di moltiplicazione e trasformazione genica.** Patent: IPO 2000A-000305
- Murashige T, Skoog F: **A revised medium for rapid growth and bioassay with tobacco tissue cultures.** *Physiologia Plantarum* 1962, **15**:473-497
- Arunnangathan K, Earle ED: **Estimating of nuclear DNA content in plants by flow cytometry.** *Plant Mol Biology Report* 1991, **7**(9):229-233
- Koncz C, Schell J: **The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of *Agrobacterium* binary vector.** *Mol Gen Genet* 1986, **204**:383-396
- Quorin M, Lepoivre P: **Improved media for in vitro culture of *Prunus* spp.** *Acta Horticulturae* 1977, **78**:437-442

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



BioMedcentral.com

Submit your manuscript here:

http://www.biomedcentral.com/manuscript/

editorial@biomedcentral.com