



## Comment on "Influence of HLA-C Expression Level on HIV Control"

Elisa Lo Monaco *et al.*  
*Science* **341**, 1175 (2013);  
DOI: 10.1126/science.1241266

*This copy is for your personal, non-commercial use only.*

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

**The following resources related to this article are available online at [www.sciencemag.org](http://www.sciencemag.org) (this information is current as of June 25, 2014 ):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/341/6151/1175.2.full.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/341/6151/1175.2.full.html#related>

This article **cites 11 articles**, 7 of which can be accessed free:

<http://www.sciencemag.org/content/341/6151/1175.2.full.html#ref-list-1>

This article has been **cited by** 1 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/341/6151/1175.2.full.html#related-urls>

This article appears in the following **subject collections**:

Medicine, Diseases

<http://www.sciencemag.org/cgi/collection/medicine>

Technical Comments

[http://www.sciencemag.org/cgi/collection/tech\\_comment](http://www.sciencemag.org/cgi/collection/tech_comment)

# Comment on “Influence of HLA-C Expression Level on HIV Control”

Elisa Lo Monaco,<sup>1</sup> Elisa Tremante,<sup>1</sup> Priscilla Biswas,<sup>2</sup> Martin P. Cranage,<sup>3</sup> Donato Zipeto,<sup>4</sup> Alberto Beretta,<sup>2</sup> Patrizio Giacomini<sup>1\*</sup>


Apps *et al.* (Reports, 5 April 2013, p. 87) found that high human leukocyte antigen C (HLA-C) expression favors HIV-1 control. However, as noted here, HLA-C was assessed with a monoclonal antibody (DT9) that cross-reacts with HLA-E. In the context of the available evidence, this is consistent with the idea that the two leukocyte antigens collaborate to keep the HIV-1 virus at bay.

A report recently published in *Science* demonstrated that the control of HIV-1 infection is positively influenced by high human leukocyte antigen C (HLA-C) expression (1). We note that this observation largely relies on extensive flow cytometry testing with a monoclonal antibody (mAb) named DT9, and draw attention to the original paper in which this mAb was described to primarily bind HLA-E, HLA-C being secondarily detected as a result of cross-reactivity (2).

The claim by Apps *et al.* for operational HLA-C specificity of mAb DT9 is based on extensive background information previously presented by the same group, particularly the lack of coexpressed HLA-E on the surface of human peripheral blood mononuclear cells (PBMCs), as assessed by a mAb named MEM-E/08 (3). However, this finding has subsequently been challenged by Corrah *et al.*, who have instead documented PBMC staining with another mAb to HLA-E, called 3D12 (4). Further adding to this controversy, and in agreement with Corrah *et al.*, we have shown that mAb MEM-E/08 reacts with poorly expressed, unstable, unfolded HLA-E molecules free of their light chain subunit ( $\beta_2m$ ). In addition, we found that HLA-E levels are often underestimated, because a fraction of HLA-E is surface-expressed as a tightly  $\beta_2m$ -associated conformer poorly reactive with the available mAbs, including to some extent even the conformational mAb 3D12 (5). Altogether, these results favor the possibility that HLA-E substantially contributes to DT9 reactivity on the surface of many different human cells. HLA-E cross-reactivity, even marginal, may not be negligible in this setting, since HLA-C is

the lowest expressed classical class I molecule [reviewed in (6)].

On this basis, we suggest that mAb DT9 detects combined HLA-C/HLA-E expression and



**Fig. 1. HLA-C and HLA-E: linked expression, reactivity with mAb DT9, and possible collaborative protection from HIV-1.** Two patterns of mAb DT9 reactivity are depicted, implying different HIV-1 control models. The first (A), proposed by Apps *et al.*, is based on a restricted HLA-C reactivity of DT9 and is consistent with demonstrated, HLA-C–specific protective effects (1, 11). The second (B), hypothesized here on the basis of the previously observed HLA-E cross-reactivity with DT9, envisages a compound multilocus effect. Because HLA-E requires peptides donated from the signal sequence of HLA-C for its stable assembly with  $\beta_2m$ , we suggest that the two class I HLA molecules are coordinately expressed: The higher HLA-C is, the higher is the surface HLA-E pool. Because they are expressed in proportional amounts, HLA-E does not prevent the appreciation of differences in HLA-C expression by mAb DT9. Evidence for a protective role of HLA-E remains so far indirect.

that this combination, rather than HLA-C alone, may be considered the most accurate readout for the outcome of HIV-1 infection. Essentially all the HLA-C alleles, but only some HLA-A and HLA-B alleles, encode signal sequences potentially acting as donors of stabilizing HLA-E ligands (7, 8) (also see [www.ebi.ac.uk/ipd/imgt/hla](http://www.ebi.ac.uk/ipd/imgt/hla)), providing direct insight into linked, and possibly coordinated, expression of HLA-C and HLA-E (Fig. 1). Coordination would result in HLA-E similarly contributing to DT9 binding in different HLA-C–expressing cells. Thus, HLA-E cross-reactivity would not preclude the discrimination of HLA-C–high from HLA-C–low alleles, a discrimination that is central to the HIV protection model put forward in several papers by the authors of (1). Protection from, and susceptibility to, HIV infection may then be interpreted in the context of a compound multilocus effect.

This alternative interpretation does not contradict, but rather complements, the original interpretation of Apps *et al.* (1). High HLA-C would incite cytotoxic T cell recognition, exactly as outlined by these authors, whereas high HLA-E may be instrumental in directing the lysis of virus-infected cells by certain HLA-E–restricted T cell subsets, as shown in the case of cytomegalovirus (9). Alternatively, or in addition, inhibitory HLA-E may engender a tolerogenic effect that paradoxically prevents the establishment of productive viral infection, as described in a macaque model of simian immunodeficiency virus vaccination (10).

## References and Notes

1. R. Apps *et al.*, *Science* **340**, 87–91 (2013).
2. V. M. Braud, D. S. J. Allan, D. Wilson, A. J. McMichael, *Curr. Biol.* **8**, 1–10 (1998).
3. R. Thomas *et al.*, *Nat. Genet.* **41**, 1290–1294 (2009).
4. T. W. Corrah *et al.*, *J. Virol.* **85**, 3367–3374 (2011).
5. E. Lo Monaco *et al.*, *J. Immunol.* **181**, 5442–5450 (2008).
6. L. Sibilio *et al.*, *J. Biol. Chem.* **283**, 1267–1274 (2008).
7. N. Lee, D. R. Goodlett, A. Ishitani, H. Marquardt, D. E. Geraghty, *J. Immunol.* **160**, 4951–4960 (1998).
8. A. G. Brooks *et al.*, *J. Immunol.* **162**, 305–313 (1999).
9. C. Romagnani *et al.*, *Hum. Immunol.* **65**, 437–445 (2004).
10. W. Lu *et al.*, *Cell Reports* **2**, 1736–1746 (2012).
11. M. E. Blais *et al.*, *J. Immunol.* **188**, 4663–4670 (2012).

**Acknowledgments:** Supported by Associazione Italiana per la Ricerca sul Cancro (AIRC), grant 40H92 (national AIDS program from Istituto Superiore di Sanità, Rome, Italy), and Europrize Network of Excellence, Life Sciences Programme, European Commission. The authors have no conflicts of interest.

30 May 2013; accepted 12 August 2013  
10.1126/science.1241266

<sup>1</sup>Istituto Nazionale Tumori Regina Elena, Via delle Messi d’Oro 156, 00158 Roma, Italy. <sup>2</sup>Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Ospedale e Istituto Scientifico San Raffaele, Via Olgettina 58, 20132 Milano, Italy. <sup>3</sup>Centre for Infection and Immunity, St George’s University of London, London, UK. <sup>4</sup>Department of Life and Reproduction Sciences, University of Verona, Strada le Grazie 8, 37134 Verona, Italy.

\*Corresponding author. E-mail: giacomini@ifo.it